

# Ideal interpolation: Mourrain's condition *vs* $D$ -invariance

C. de Boor

By definition (see [Bi]), **ideal interpolation** is provided by a linear projector whose kernel is an ideal in the ring  $\Pi$  of polynomials (in  $d$  real ( $\mathbb{F} = \mathbb{R}$ ) or complex ( $\mathbb{F} = \mathbb{C}$ ) variables). The standard example is Lagrange interpolation; the most general example has been called ‘Hermite interpolation’ (in [M] and [citB05]) since that is what it reduces to in the univariate case.

Ideal projectors also occur in computer algebra, as the maps that associate a polynomial with its *normal form* with respect to an ideal; see, e.g., [CLO]. It is in this latter context that Mourrain [Mo] poses and solves the following problem. *Among all linear projectors  $N$  on*

$$\Pi_1(F) := \sum_{j=0}^d ()_j F$$

*with range the linear space  $F$ , characterize those that are the restriction to  $\Pi_1(F)$  of an ideal projector with range  $F$ .* Here,

$$()_j := ()^{\varepsilon_j}, \quad \varepsilon_j := (\delta_{jk} : k = 1:d), \quad j = 0:d,$$

with

$$()^\alpha : \mathbb{F}^d \rightarrow \mathbb{F} : x \mapsto x^\alpha := \prod_{j=1}^d x(j)^{\alpha(j)}$$

a handy if nonstandard notation for the **monomial with exponent**  $\alpha \in \mathbb{Z}_+^d$ . I also use the corresponding notation

$$D_j$$

for the derivative with respect to the  $j$ th argument, and

$$D^\alpha := \prod_{j=1}^d D_j^{\alpha(j)}, \quad \alpha \in \mathbb{Z}_+^d.$$

To state Mourrain's result, I also need the following, standard, notations. The (total) **degree** of the polynomial  $p \neq 0$  is the nonnegative integer

$$\deg p := \max\{|\alpha| : \widehat{p}(\alpha) \neq 0\},$$

with

$$p =: \sum_{\alpha} ()^\alpha \widehat{p}(\alpha),$$

and

$$|\alpha| := \sum_j \alpha(j),$$

while

$$\Pi_{<n} := \{p \in \Pi : \deg p < n\}.$$

**Theorem 1 ([Mo]).** *Let  $F$  be a finite-dimensional linear subspace of  $\Pi$  satisfying **Mourrain's condition**:*

$$(2) \quad f \in F \implies f \in \Pi_1(F \cap \Pi_{<\deg f}),$$

and let  $N$  be a linear projector on  $\Pi_1(F)$  with range  $F$ . Then, the following are equivalent:

- (a)  $N$  is the restriction to  $\Pi_1(F)$  of an ideal projector with range  $F$ .
- (b) The linear maps  $M_j : F \rightarrow F : f \mapsto N((\cdot)_j f)$ ,  $j = 1:d$ , commute.

For a second proof of this theorem and some unexpected use of it in the setting of ideal interpolation, see [citB05].

Mourrain's condition (2) implies that, if  $F$  contains an element of degree  $k$ , it must also contain an element of degree  $k - 1$ . In particular, if  $F$  is nontrivial, then it must contain a constant polynomial. This explains why Mourrain [Mo] calls a linear subspace satisfying his condition **connected to 1**. Since the same argument can be made in case  $F$  is  $D$ -invariant, this raises the question what connection if any there might be between these two properties.

In particular, for the special case  $d = 1$ , if  $F$  is a linear subspace of dimension  $n$  and either satisfying Mourrain's condition or being  $D$ -invariant, then, necessarily,  $F = \Pi_{<n}$ . More generally, if  $F$  is an  $n$ -dimensional subspace in the subring generated by

$$\langle \cdot, y \rangle : \mathbb{F}^d \rightarrow \mathbb{F} : x \mapsto \langle x, y \rangle := \sum_{j=1}^d x(j)y(j)$$

for some  $y \neq 0$ , then, either way,

$$F = \text{ran}[\langle \cdot, y \rangle^{j-1} : j = 1:n] := \left\{ \sum_{j=1}^n \langle \cdot, y \rangle^{j-1} a(j) : a \in \mathbb{F}^n \right\}.$$

As a next example, assume that  $F$  a **monomial** space (meaning that it is spanned by monomials). If such  $F$  is  $D$ -invariant, then, with each  $(\cdot)^\alpha$  for which  $\alpha - \varepsilon_j \in \mathbb{Z}_+^d$ , it also contains  $(\cdot)^{\alpha - \varepsilon_j}$  and therefore evidently satisfies Mourrain's condition.

Slightly more generally, assume that  $F$  is **dilation-invariant**, meaning that it contains  $f(h \cdot)$  for every  $h > 0$  if it contains  $f$  or, equivalently,  $F$  is spanned by homogeneous polynomials. Then every  $f \in F$  is of the form

$$f =: f_\uparrow + f_0,$$

with  $f_\uparrow$  the **leading** term of  $f$ , i.e., the unique homogeneous polynomial for which

$$\deg(f - f_\uparrow) < \deg f,$$

hence in  $F$  by dilation-invariance, therefore also

$$f_0 \in F_{<\deg f} := F \cap \Pi_{<\deg f},$$

while, by the homogeneity of  $f_\uparrow$ ,

$$\sum_{j=1}^d ()_j D_j(f_\uparrow) = (\deg f) f_\uparrow$$

(this is **Euler's theorem for homogeneous functions**; see, e.g., [Encycl: p281] which gives the reference [E: §225 on p154]). If now  $F$  is also  $D$ -invariant, then  $D_j(f_\uparrow) \in F_{<\deg f}$ , hence, altogether,

$$f \in \Pi_1(F_{<\deg f}), \quad f \in F.$$

In other words, *if a dilation-invariant finite-dimensional subspace  $F$  of  $\Pi$  is  $D$ -invariant, then it also satisfies Mourrain's condition.*

On the other hand, the linear space

$$\text{ran}[()^0, ()^{1,0}, ()^{1,1}]$$

fails to be  $D$ -invariant even though it satisfies Mourrain's condition and is monomial, hence dilation-invariant.

The final example, of a space that is  $D$ -invariant but does not satisfy Mourrain's condition, is slightly more complicated. In its discussion, I find it convenient to refer to

$$\text{supp } \widehat{p}$$

as the '**support**' of the polynomial  $p = \sum_{\alpha} ()^{\alpha} \widehat{p}(\alpha)$ , with the quotation marks indicating that it isn't actually the support of  $p$  but, rather, the support of its coefficient sequence,  $\widehat{p}$ . The example is provided by the  $D$ -invariant space  $F$  generated by the polynomial

$$p = ()^{1,7} + ()^{3,3} + ()^{5,0},$$

hence the 'support' of  $p$  is

$$\text{supp } \widehat{p} = \{(1, 7), (3, 3), (5, 0)\}$$

(see (4) below). Here are a first few elements of  $F$ :

$$D_1 p = ()^{0,7} + 3()^{2,3} + 5()^{4,0}, \quad D_2 p = 7()^{1,6} + 3()^{3,2},$$

hence

$$D_1 D_2 p = 7()^{0,6} + 9()^{2,2}, \quad D_2^2 p = 42()^{1,5} + 6()^{3,1},$$

also

$$D_1^2 p = 6()^{1,3} + 20()^{3,0}, \quad D_1 D_2^2 p = 42()^{0,5} + 18()^{2,1},$$

etc. This shows (see (4) below) that any  $q \in \Pi_1(F_{<\deg p})$  having some ‘support’ in  $\text{supp } \widehat{p}$  is necessarily a weighted sum of  $(\ )_1 D_1 p$  and  $(\ )_2 D_2 p$  (and, perhaps, others not having any ‘support’ in  $\text{supp } \widehat{p}$ ), yet  $(p, (\ )_1 D_1 p, (\ )_2 D_2 p)$  is linearly independent ‘on’  $\text{supp } \widehat{p}$ , as the matrix

$$\begin{bmatrix} 1 & 1 & 7 \\ 1 & 3 & 3 \\ 1 & 5 & 0 \end{bmatrix}$$

(of their coefficients indexed by  $\alpha \in \text{supp } \widehat{p}$ ) is evidently 1-1. Consequently,  $p \notin \Pi_1(F_{<\deg p})$ , i.e., this  $F$  does not satisfy Mourrain’s condition.

This space also provides the proof that, in Theorem 1, one may not, in general, replace Mourrain’s condition by  $D$ -invariance.

**Proposition 3.** *Let  $F$  be the  $D$ -invariant space spanned by*

$$p = (\ )^{1,7} + (\ )^{3,3} + (\ )^{5,0}.$$

*Then there exists a linear projector,  $N$ , on  $\Pi_1(F)$  with range  $F$  for which (b) but not (a) of Theorem 1 is satisfied.*

**Proof:** For  $\alpha, \beta \in \mathbb{Z}_+^d$ , set

$$[\alpha \dots \beta] := \{\gamma \in \mathbb{Z}_+^d : \alpha \leq \gamma \leq \beta\},$$

with

$$\alpha \leq \gamma := \forall j \quad \alpha(j) \leq \gamma(j).$$

With this, we determine a basis for  $F$  as follows.

Since  $D^{0,4}p$  is a positive scalar multiple of  $(\ )^{1,3}$ , we know, by the  $D$ -invariance of  $F$ , that

$$\{(\ )^\zeta : \zeta \in [(0,0) \dots (1,3)]\} \subset F.$$

This implies, considering  $D^{2,0}p$ , that  $(\ )^{3,0}$ , hence also  $(\ )^{2,0}$ , is in  $F$ . Hence, altogether,

$$F = \Pi_{\Xi_0} \oplus \text{ran}[D^\alpha p : \alpha \in [(0,0) \dots (1,3)]],$$

with

$$\Pi_\Gamma := \text{ran}[(\ )^\gamma : \gamma \in \Gamma]$$

and

$$\Xi_0 := [(0,0) \dots (1,3)] \cup \{(2,0), (3,0)\}.$$

This provides the convenient basis

$$b_\Xi := [b_\xi : \xi \in \Xi]$$

for  $F$ , indexed by

$$\Xi := \Xi_0 \cup \Xi_1, \quad \Xi_1 := [(0,4) \dots (1,7)],$$

namely

$$b_\xi := \begin{cases} ()^\xi, & \xi \in \Xi_0; \\ D^{(1,7)-\xi}p, & \xi \in \Xi_1. \end{cases}$$

The following schema indicates the sets  $\text{supp } \widehat{p}$ ,  $\Xi_0$ , and  $\Xi_1$ , as well as the sets  $\partial\Xi_0$  and  $\partial\Xi_1$  defined below:

$$(4) \quad \begin{array}{ccccccc} & \times & \times & & & & \otimes : \text{supp } \widehat{p} \\ & 1 & \oplus & \times & & & 0 : \Xi_0 \\ & 1 & 1 & \times & & & 1 : \Xi_1 \\ & 1 & 1 & \times & & & + : \partial\Xi_0 \\ & 1 & 1 & \times & & & \times : \partial\Xi_1 \\ & 0 & 0 & + & \otimes & & \\ & 0 & 0 & + & & & \\ & 0 & 0 & + & + & & \\ & 0 & 0 & 0 & 0 & + & \otimes \end{array}$$

Now, let  $N$  be the linear projector on  $\Pi_1(F)$  with range  $F$  and kernel  $\text{ran}[b_Z]$ , with  $b_Z$  obtained by thinning

$$[b_\Xi, ()_1b_\Xi, ()_2b_\Xi]$$

to a basis  $[b_\Xi, b_Z]$  for  $\Pi_1(F)$ . This keeps the maps  $M_j : F \rightarrow F : f \mapsto N(()_j f)$  very simple since, as we shall see, for many of the  $\xi \in \Xi$ ,  $()_j b_\xi$  is an element of the extended basis  $[b_\Xi, b_Z]$ , hence  $N$  either reproduces it or annihilates it.

Specifically, it is evident that the following are in  $F$ , hence not part of  $b_Z$ :

$$()_1 b_\xi, \quad \xi \in [(0,0) \dots (0,2)],$$

$$()_2 b_\xi, \quad \xi \in [(0,0) \dots (1,3)].$$

Further, for each

$$\zeta \in \partial\Xi_0 \cup \partial\Xi_1,$$

with

$$\partial\Xi_0 := \{(2,3), (2,2), (2,1), (3,1), (4,0)\}, \quad \partial\Xi_1 = \{[(2,4) \dots (2,7)], (1,8), (0,8)\},$$

there is  $\xi \in \Xi$  so that, for some  $j$ ,

$$\zeta - \xi = \varepsilon_j := \begin{cases} (1,0), & j = 1; \\ (0,1), & j = 2. \end{cases}$$

Set, correspondingly,

$$b_\zeta := ()_j b_\xi.$$

Then, none of these is in  $F$ , and, among them, each  $b_\zeta$  is the only one having some ‘support’ at  $\zeta$ , hence they form a linearly independent sequence. Therefore, each such  $b_\zeta$  is in  $b_Z$ .

The remaining candidates for membership in  $b_Z$  require a more detailed analysis. We start from the ‘top’, showing also along the way that (b) of Theorem 1 holds for this  $F$  and  $N$  by verifying that

$$(5) \quad M_1 M_2 = M_2 M_1 \text{ on } b_\xi$$

for every  $\xi \in \Xi$ .

$\xi = (1, 7)$ : As already pointed out, both  $(\ )_1 b_{1,7}$  and  $(\ )_2 b_{1,7}$  are in  $b_Z$ , hence (5) holds trivially for  $\xi = (1, 7)$ .

$\xi = (0, 7), (1, 6)$ : Both  $(\ )_1 b_{0,7} = (\ )^{1,7} + 3(\ )^{3,3} + 5(\ )^{5,0}$  and  $(\ )_2 b_{1,6} = 7(\ )^{1,7} + 3(\ )^{3,3}$  have their ‘support’ in that of  $p = b_{1,7} = (\ )^{1,7} + (\ )^{3,3} + (\ )^{5,0}$ , while, as pointed out and used earlier, the three are independent. Hence  $(\ )_1 b_{0,7}, (\ )_2 b_{1,6} \in b_Z$ , while we already pointed out that  $(\ )_2 b_{0,7}, (\ )_1 b_{1,6} \in b_Z$ , therefore (5) holds trivially.

$\xi = (0, 6), (1, 5)$ : Both  $(\ )_1 b_{0,6} = 7(\ )^{1,6} + 9(\ )^{3,2}$  and  $(\ )_2 b_{1,5} = 42(\ )^{1,6} + 6(\ )^{3,2}$  have their ‘support’ in that of  $b_{1,6} = 7(\ )^{1,6} + 3(\ )^{3,2}$ , but neither is a scalar multiple of  $b_{1,6}$ . Hence, one is in  $b_Z$  and the other is not. Which is which depends on the ordering of the columns of  $[b_\Xi, (\ )_1 b_\Xi, (\ )_2 b_\Xi]$ . Assume the ordering such that  $(\ )_2 b_{1,5} \in b_Z$ . Then, since we already know that  $(\ )_1 b_{1,5} \in b_Z$ , (5) holds trivially for  $\xi = (1, 5)$ . Further,  $(\ )_1 b_{0,6} = 4b_{1,6} - (1/2)(\ )_2 b_{1,5}$ , hence  $M_1 b_{0,6} = 4b_{1,6}$ , while we already know that  $(\ )_2 b_{1,6} \in b_Z$  therefore,  $M_2 M_1 b_{0,6} = 0$ . On the other hand,  $(\ )_2 b_{0,6} = 7(\ )^{0,7} + 3(\ )^{3,3}$  has its ‘support’ in that of  $b_{0,7} = (\ )^{0,7} + 3(\ )^{3,3} + 5(\ )^{4,0}$  but is not a scalar multiple of it, hence is in  $b_Z$ , and therefore already  $M_2 b_{0,6} = 0$ . Thus, (5) also holds for  $\xi = (0, 6)$ .

$\xi = (0, 5), (1, 4)$ : Both  $(\ )_1 b_{0,5} = 42(\ )^{1,5} + 18(\ )^{3,1}$  and  $(\ )_2 b_{1,4} = 210(\ )^{1,5} + 6(\ )^{3,1}$  have their ‘support’ in that of  $b_{1,5} = 42(\ )^{1,5} + 6(\ )^{3,1}$  but  $(\ )^{3,1} = b_{3,1}$  was already identified as an element of  $b_Z$ , hence neither  $(\ )_1 b_{0,5}$  nor  $(\ )_2 b_{1,4}$  is in  $b_Z$ . But, since  $(\ )^{3,1} \in b_Z$ , and so  $b_{1,5} = N b_{1,5} = N(42(\ )^{1,5})$ , we have  $M_1 b_{0,5} = b_{1,5}$  and  $M_2 b_{1,4} = 5b_{1,5}$ . Since we already know that  $(\ )_1 b_{1,5} \in b_Z$ , it follows that  $M_1 M_2 b_{1,4} = 0$  while we already know that  $(\ )_1 b_{1,4} \in b_Z$ , hence already  $M_1 b_{1,4} = 0$ . Therefore, (5) holds for  $\xi = (1, 4)$ . Further, we already know that  $(\ )_2 b_{1,5} \in b_Z$ , hence  $M_2 M_1 b_{0,5} = 0$ , while  $(\ )_2 b_{0,5} = 42(\ )^{0,6} + 18(\ )^{2,2}$  has the same ‘support’ as  $b_{0,6} = 7(\ )^{0,6} + 9(\ )^{2,2}$  but is not a scalar multiple of it, hence is in  $b_Z$  and, therefore, already  $M_2 b_{0,5} = 0$ , showing that (5) holds for  $\xi = (0, 5)$ .

$\xi = (0, 4)$ :  $(\ )_2 b_{0,4} = 210(\ )^{0,5} + 18(\ )^{2,1} = 5b_{0,5} - 72b_{2,1}$ , with  $b_{2,1} \in b_Z$ , hence  $(\ )_2 b_{0,4}$  is not in  $b_Z$  and  $M_2 b_{0,4} = 5b_{0,5}$ , therefore  $M_1 M_2 b_{0,4} = 5M_1 b_{0,5} = 5b_{1,5}$ , the last equation from the preceding paragraph. On the other hand,  $(\ )_1 b_{0,4} = 210(\ )^{1,4} + 18(\ )^{3,0} = b_{1,4} + 12b_{3,0}$ , with both  $b_{1,4}$  and  $b_{3,0}$  in  $F$ , hence  $(\ )_1 b_{0,4}$  is not in  $b_Z$ , and  $M_1 b_{0,4} = b_{1,4} + 12b_{3,0}$ , therefore, since  $(\ )_2 b_{3,0} = b_{3,1} \in b_Z$ ,  $M_2 M_1 b_{0,4} = M_2 b_{1,4} = 5b_{1,5}$ , the last equation from the preceding paragraph. Thus, (5) holds for  $\xi = (0, 4)$ .

$\xi = (1, 3)$ : We already know that  $(\ )_1 b_{1,3} = b_{2,3} \in b_Z$  and therefore already  $M_1 b_{1,3} = 0$ , while  $(\ )_2 b_{1,3} = (\ )^{1,4} = (b_{1,4} - 6b_{3,0})/210 \in F$ , therefore  $210M_1 M_2 b_{1,3} = M_1 b_{1,4} = 0$ , thus (5) holds for  $\xi = (1, 3)$ .

For the remaining  $\xi \in \Xi$ , each  $b_\xi$  is a monomial, hence  $(\ )_j b_\xi$  is again a monomial, and either in  $F$  or not and, if not, then its exponent is in

$$\partial\Xi_0 := \{(2, 3), (2, 2), (2, 1), (3, 1), (4, 0)\}.$$

Moreover,  $(\ )_1(\ )_2 b_\xi$  is in  $F$  iff  $(\ )_2(\ )_1 b_\xi$  is. Hence, (5) also holds for the remaining  $\xi \in \Xi$ . This finishes the proof that, for this  $F$  and  $N$ , (b) of Theorem 1 holds.

It remains to show that, nevertheless, (a) of Theorem 1 does not hold. For this, observe that  $(\ )^{2,1}$  and  $(\ )^{4,0}$  are in  $\ker N$ , as is, e.g.,  $(\ )_2 b_{1,6} = 7(\ )^{1,7} + 3(\ )^{3,3}$ , hence  $p = (\ )^{1,7} + (\ )^{3,3} + (\ )^{5,0}$  is in the ideal generated by  $\ker N$ , making it impossible for  $N$  to be the restriction to  $\Pi_1(F)$  of an ideal projector  $P$  with range  $F$  since this would place the nontrivial  $p$  in both  $\ker P$  and  $\text{ran } P$ .  $\square$

#### References

- [Bi] G. Birkhoff (1979), “The algebra of multivariate interpolation”, in *Constructive approaches to mathematical models* (C. V. Coffman and G. J. Fix, eds), Academic Press (New York), 345–363.
- [B] C. de Boor (1979), “Ideal interpolation”, in **the specified proceedings does not exist in our files** (■■■■, ed), (New York), xxx–xxx.
- [CLO] David Cox, John Little, and Donal O’Shea (1992), *Ideals, Varieties, and Algorithms*, Undergraduate Texts in Math., Springer-Verlag (New York).
- [Encycl] (1898–1904), *Encycl. mathem. Wissenschaften, Erster Band*, Teubner (Leipzig).
- [E] L. Euler (1787), *Calculus differentialis 1*, Ticini (Italy).
- [M] H. M. Möller (1976), “Mehrdimensionale Hermite-Interpolation und numerische Integration”, *Math. Z.* **148**, 107–118.
- [Mo] B. Mourrain (1999), “A new criterion for normal form algorithms”, in *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, 13th Intern. Symp., AAECC-13, Honolulu, Hawaii USA, Nov. ’99, Proc.* (Mark Fossorier, Hideki Imai, Shu Lin, Alan Pol, eds), Springer Lecture Notes in Computer Science, 1719, Springer-Verlag (Heidelberg), 430–443.