

Best Approximation Properties of Spline Functions of Odd Degree

CARL DE BOOR

Communicated by GARRETT BIRKHOFF

Introduction. In [4], interpolation by cubic spline functions is discussed, and some best approximation properties of the cubic spline fit are described. This note extends the results of [4], in a somewhat modified form, to spline functions of odd degree $m = 2k - 1$, $k \geq 2$.

Definition. A spline function of degree m with joints $\xi_1 < \xi_2 < \dots < \xi_n$ is defined as a function $F(x)$ with the following two properties ([2], p. 67):

- a. In each of the intervals $(-\infty, \xi_1)$, $[\xi_1, \xi_2)$, \dots , $[\xi_n, \infty)$, $F(x)$ is a polynomial of degree m ;
- b. $F(x)$ has continuous derivatives through the $(m - 1)^{\text{st}}$, or, for short, $F(x) \in C^{m-1}$.

The class of functions $F(x)$ with these properties will be denoted by $S_m(\xi_1, \dots, \xi_n)$.

The following lemma establishes the existence and uniqueness of a spline function of degree $(2k - 1)$ with $(n - 1)$ joints which coincides with a given function $f(x)$ at $(n + 1)$ prescribed points. The lemma is a consequence of Theorem 2 in [3], p. 258.

Lemma 1. Let $f(x)$ be any function of class $C^k[a, b]$. For each choice of $n + 1$ abscissae x_i , $a = x_0 < x_1 < \dots < x_n = b$, there exists exactly one spline function in $S_{2k-1}(x_1, \dots, x_{n-1})$, denoted by $\bar{s}(x)$, such that

- (1) $\bar{s}(x_i) = f(x_i), \quad i = 0, \dots, n,$
- (2) $\bar{s}^{(k+i)}(x_i) = 0, \quad i = 0, n; \quad j = 0, \dots, k - 2,$

where $\bar{s}^{(m)}(x)$ denotes the m^{th} derivative of $\bar{s}(x)$.

It has been known for some time (cf., e.g. [2], p. 67) that in the case $k = 2$ of cubic splines the interpolating function $\bar{s}(x)$ minimizes $\int_a^b [u''(x)]^2 dx$ among all functions $u(x) \in C^2$ which coincide with $f(x)$ at the points x_i , $i = 0, \dots, n$. The

cubic spline function $\bar{s}(x)$ gives therefore, approximately, the shape of a thin beam or "spline", which is forced to go through the points $\{x_i, f(x_i)\}$, $i = 0, \dots, n$. This result can be seen by considering the integral $\int [u''(x)]^2 dx$ as a linearized approximation to the strain energy of a thin beam, which is $\int u''^2/(1 + u'^2)^{5/2} dx$. Thus $\bar{s}(x)$ minimizes the strain energy subject to the geometrical constraints stated (cf. [1], p. 92-98). The corresponding nonlinear problem was first considered by L. Euler and D. Bernoulli.

The inner product

$$(3) \quad (f, g)_k = \int_a^b f^{(k)}(x)g^{(k)}(x) dx$$

is defined for any two functions f, g which have square-integrable k^{th} derivatives on $[a, b]$. It defines a pseudo-norm

$$(4) \quad \|f\|_k = [(f, f)_k]^{1/2}$$

on the linear space $C^k[a, b]$, in which $\|f\|_k = 0$ if and only if $f(x)$ is a polynomial of degree $(k - 1)$ or less.

Theorem 1. Among all the functions $u(x) \in C^k[a, b]$, which satisfy (1') $u(x_i) = f(x_i)$, $i = 0, \dots, n$, the pseudo-norm $\|u\|_k$ is minimized by $\bar{s}(x)$.

In this sense, the spline function $\bar{s}(x)$ is the smoothest function interpolating $f(x)$ at the points x_i , $i = 0, \dots, n$.

This theorem is a direct consequence of the following lemma.

Lemma 2. If $f(x) \in C^k[a, b]$, and $\bar{s}(x) \in S_{2k-1}(x_0, \dots, x_n)$ satisfies (1) and (2), then

$$(5) \quad \|f\|_k^2 - \|\bar{s}\|_k^2 = \|f - \bar{s}\|_k^2.$$

Proof. Let $\eta(x) \equiv f(x) - \bar{s}(x)$. The right-hand side of (5) may be written as

$$(6) \quad \|\eta\|_k^2 = \|f\|_k^2 - \|\bar{s}\|_k^2 - 2(\eta, \bar{s})_k.$$

By successive integration by parts, one has

$$(7) \quad (\eta, \bar{s})_k = \left[\sum_{i=0}^{k-2} (-1)^i \eta^{(k-1-i)}(x) \bar{s}^{(k+i)}(x) \right] \Big|_a^b + (-1)^{k-1} \int_a^b \eta'(x) \bar{s}^{(2k-1)}(x) dx.$$

The first term of the right-hand side of (7) vanishes because of (2). Since $\bar{s}^{(2k-1)}(x)$ is a constant in each of the intervals (x_i, x_{i+1}) , $i = 0, \dots, n-1$, ($x_0 = a$, $x_n = b$), one has for the second term

$$(8) \quad \int_a^b \eta'(x) \bar{s}^{(2k-1)}(x) dx = \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \eta'(x) \bar{s}^{(2k-1)}(x) dx \\ = \sum_{i=0}^{n-1} [\eta(x_{i+1}) - \eta(x_i)] \bar{s}^{(2k-1)}(i\text{-th interval}),$$

and this, by (1), is zero.

Remark. Lemmas 1 and 2 remain true, if condition (2) is replaced by

$$(2') \quad \bar{s}^{(i)}(x_i) = f^{(i)}(x_i), \quad i = 0, n; \quad j = 1, \dots, k - 1.$$

Lemma 2'. If $f(x) \in C^k[a, b]$, and $\bar{s}(x) \in S_{2k-1}(x_1, \dots, x_{n-1})$ satisfies (1) and (2') then

$$(5) \quad \|f\|_k^2 - \|\bar{s}\|_k^2 = \|f - \bar{s}\|_k^2.$$

For the remainder of this note, let $\bar{s}(x) \in S_{2k-1}(x_1, \dots, x_{n-1})$ denote the unique spline function of degree $2k - 1$ which satisfies (1) and (2'), and hence (5).

Theorem 1'. Among the functions $u(x) \in C^k[a, b]$, which satisfy (1) and (2'), (with $\bar{s}(x)$ replaced by $u(x)$), the norm $\|u\|_k$ is minimum for $\bar{s}(x)$.

Lemma 2' not only implies Theorem 1', but provides a characterization of the best approximation $s^*(x)$ to $f(x) \in C^k[a, b]$ by spline functions $s(x) \in S_{2k-1}(x_1, \dots, x_{n-1})$ with respect to the measure of approximation

$$(9) \quad \|f - s\|_k.$$

A best approximation $s^*(x) \in S_{2k-1}(x_1, \dots, x_{n-1})$ has to satisfy

$$(10) \quad \|f - s^*\|_k \leq \|f - s\|_k, \quad \text{for all } s \in S_{2k-1}(x_1, \dots, x_{n-1}).$$

Since $\|f\|_k = 0$ if and only if $f(x)$ is a polynomial of degree $(k - 1)$, i.e., $f(x) \equiv P_{k-1}(x)$, best approximations are not unique; $(s(x) + P_{k-1}(x))$ is a best approximation, if $s(x)$ is.

Theorem 2. For $f(x) \in C^k[a, b]$,

$$(11) \quad s^*(x) = \bar{s}(x) + P_{k-1}(x),$$

i.e., the spline function $\bar{s}(x)$ interpolating $f(x)$ at the points $x_i, i = 0, \dots, n$, and satisfying (2'), is a best approximation to $f(x)$ by spline functions in $S_{2k-1}(x_1, \dots, x_{n-1})$ with respect to the measure of approximation (9).

Proof. Let $s(x)$ be any function in $S_{2k-1}(x_1, \dots, x_{n-1})$. In Lemma 2', replace $f(x)$ by $(f(x) - s(x))$. Then $(\bar{s}(x) - s(x))$ is the corresponding unique function in $S_{2k-1}(x_1, \dots, x_{n-1})$ satisfying (1) and (2'), so that

$$(12) \quad \|f - \bar{s}\|_k^2 = \|f - s\|_k^2 - \|\bar{s} - s\|_k^2.$$

Hence

$$\|f - \bar{s}\|_k \leq \|f - s\|_k,$$

with equality holding if and only if $\|\bar{s} - s\|_k = 0$,—i.e., when $s(x) = \bar{s}(x) + P_{k-1}(x)$.

REFERENCES

- [1] T. L. SYNGE & B. A. GRIFFITH, *Principles of mechanics*, McGraw-Hill (1949).
- [2] I. J. SCHOENBERG, Approximation of equidistant data, *Quart. Appl. Math.*, **4** (1946) 45-99.
- [3] I. J. SCHOENBERG & A. WHITNEY, On Polya frequency functions, III, *Trans. A. M. S.*, **74** (1953) 246-259.
- [4] J. L. WALSH, J. H. AHLBERG & E. N. NILSON, Best approximation properties of the spline fit, *J. Math. Mech.*, **11** (1962) 225-234.

General Motors Research
Laboratories