

The Numerically Stable Reconstruction of a Jacobi Matrix from Spectral Data

C. de Boor*

*Mathematics Research Center
University of Wisconsin—Madison
Madison, Wisconsin 53706*

and

G. H. Golub†

*Computer Science Department
Stanford University
Stanford, California 94305*

Submitted by Hans Schneider

ABSTRACT

We show how to construct, from certain spectral data, a discrete inner product for which the associated sequence of monic orthogonal polynomials coincides with the sequence of appropriately normalized characteristic polynomials of the left principal submatrices of the Jacobi matrix. The generation of these orthogonal polynomials via their three-term recurrence relation, as popularized by Forsythe, then provides a stable means of computing the entries of the Jacobi matrix. Our construction provides, incidentally, very simple proofs of known results concerning the existence and uniqueness of a Jacobi matrix satisfying given spectral data and its continuous dependence on those data.

1. INTRODUCTION

Gantmacher and Krein [3] take the term “Jacobi matrix” to mean nothing more than “tridiagonal matrix.” But it seems to have become accepted in papers on the problem of concern here to mean by “Jacobi matrix” *a real, symmetric, tridiagonal matrix whose next-to-diagonal elements are positive*. We follow this usage here, and write such a Jacobi matrix

*Sponsored by the United States Army under Contract No. DAAG29-75-C-0024.

†Supported, in part, by MRC under Army Contract No. DAAG29-75-C-0024, and in part by ERDA grant AT(04-3)-326PA #30.

J of order n as

$$J = \begin{bmatrix} a_1 & b_1 & & & & & \\ b_1 & a_2 & b_2 & & & & \\ & b_2 & a_3 & b_3 & & & \\ & & \ddots & \ddots & \ddots & & \\ & & & b_{n-2} & a_{n-1} & b_{n-1} & \\ & & & & b_{n-1} & a_n & \end{bmatrix}, \quad b_i > 0, \quad \text{all } i. \quad (1)$$

Further, we denote its left principal submatrix of order r by J_r .
 We consider the following inverse problem.

PROBLEM A. *Given the sequences $\lambda := (\lambda_i)_1^n$ and $\mu := (\mu_i)_1^{n-1}$ with*

$$\lambda_i < \mu_i < \lambda_{i+1}, \quad i = 1, \dots, n-1, \quad (S)$$

construct an n th-order Jacobi matrix J which has $\lambda_1, \dots, \lambda_n$ as its eigenvalues and μ_1, \dots, μ_{n-1} as the eigenvalues of its left principal submatrix J_{n-1} of order $n-1$.

It is well known that the eigenvalues of J_{n-1} strictly separate those of $J_n = J$, so that the condition (S) is necessary for the existence of a solution. Hochstadt [7] proved that the problem has at most one solution and proved in [8] that this solution (if it exists) depends continuously on λ and μ . L. J. Gray and D. G. Wilson [5] showed it to have at least one solution, as did O. H. Hald [6]. The latter also demonstrated in more explicit detail the continuous dependence of J on λ and μ and described an algorithm for the construction of J which, however, fails to be stable. He also announced an iterative, linearly convergent procedure for the determination of J . A different iterative procedure was developed by Barcilon [1].

By contrast, the algorithm described below in Sec. 4 is direct, i.e., not iterative, and is stable. Its derivation provides simple proofs of the results concerning Problem A just mentioned.

We also consider the following related problems.

PROBLEM B. *Given two strictly increasing sequences $\lambda := (\lambda_i)_1^n$ and $\lambda^* := (\lambda_i^*)_1^n$ with $\lambda_i < \lambda_i^* < \lambda_{i+1}$, all i , determine an n th-order Jacobi matrix J which has $\lambda_1, \dots, \lambda_n$ as its eigenvalues and for which the matrix J^* , obtained from J by changing a_n to a_n^* , has $\lambda_1^*, \dots, \lambda_n^*$ as its eigenvalues.*

PROBLEM C. *Given the strictly increasing sequence $\lambda := (\lambda_i)_1^n$, construct an n th-order persymmetric Jacobi matrix J having $\lambda_1, \dots, \lambda_n$ as its eigenvalues.*

Here, a matrix $A=(a_{ij})$ is called *persymmetric* if it is symmetric with respect to its second diagonal, i.e., if $a_{ij}=a_{n+1-j,n+1-i}$, all i and j . The Jacobi matrix (1) is persymmetric iff $a_i=a_{n+1-i}$ and $b_i=b_{n-i}$, all i .

Hochstadt [7] showed Problem C to have at most one solution. Hald [6] showed it to have at least one solution and showed the solution to depend continuously on λ .

In the analysis of these problems, the intimate connection between Jacobi matrices and orthogonal polynomials plays an essential role. We recall the salient facts of this connection in the next section.

2. JACOBI MATRICES AND ORTHOGONAL POLYNOMIALS

We continue to use the notation J_i for the left principal submatrix of order i of the Jacobi matrix (1). Let

$$p_i(t) := \det(t - J_i), \quad i = 1, \dots, n.$$

Then p_i is a monic polynomial of degree i , all i , and one verifies easily that the sequence (p_i) satisfies the three-term recurrence

$$\begin{aligned} p_i(t) &= (t - a_i) p_{i-1}(t) - b_{i-1}^2 p_{i-2}(t), & i = 1, \dots, n \\ p_{-1}(t) &= 0, & p_0(t) = 1. \end{aligned} \tag{2}$$

Conversely, if we start with a sequence (p_i) of monic polynomials with $\deg p_i = i$, all i , which also satisfies the recurrence (2), then the Jacobi matrix (1) belongs to it in the sense that then $p_i(t) = \det(t - J_i)$ for $i = 1, \dots, n$. Since the zeros of p_i are the eigenvalues of J_i , all i , we can therefore phrase Problem A equivalently as follows.

PROBLEM A'. *Given the sequences $\lambda := (\lambda_i)_1^n$ and $\mu := (\mu_i)_1^{n-1}$ with $\lambda_i < \mu_i < \lambda_{i+1}$, all i , construct sequences $a := (a_i)_1^n$ and $b := (b_i)_1^{n-1}$ so that the sequence (p_i) of polynomials given by (2) satisfies*

$$p_{n-1}(t) = \prod_{j=1}^{n-1} (t - \mu_j) \quad \text{and} \quad p_n(t) = \prod_{j=1}^n (t - \lambda_j).$$

It is clear that this problem has at most one solution, since we can always run the recurrence (2) *backwards*: If we already know the monic polynomials p_i and p_{i-1} (of degree i and $i-1$, respectively), then a_i is uniquely de-

terminated by the requirement that

$$q(t) := p_i(t) - (t - \alpha_i) p_{i-1}(t)$$

be a polynomial of degree $i - 2$. Further, the number $-\beta_{i-1}^2$ is then found as the leading coefficient of q , and p_{i-2} is then constructed by dividing q by its leading coefficient.

This construction of (p_i) satisfying (2) from p_{n-1} and p_n goes back to Wendroff [9] and has been used by Hald to solve Problem A or A' numerically. We, too, tried it in some examples and found it to be badly unstable. But, in trying to understand Hochstadt's procedure for the reconstruction of J from λ and μ [8], it occurred to us that it should be possible to construct a discrete inner product whose corresponding orthogonal polynomials satisfy (2), thus allowing us to generate a and b in the manner advocated by Forsythe [2].

We recall the details. Denote by \mathbf{P}_k the linear space of polynomials of order k , i.e., of degree $< k$, with real coefficients, and let $\langle \cdot, \cdot \rangle$ be a symmetric bilinear form which is an inner product on \mathbf{P}_n . Then there exists exactly one sequence $(q_i)_0^n$ of monic polynomials, with q_i of degree i , all i , which is orthogonal with respect to the inner product $\langle \cdot, \cdot \rangle$, i.e., for which

$$\langle q_i, q_j \rangle = 0 \quad \text{for } i \neq j.$$

One may determine q_i as the error in the best approximation from \mathbf{P}_i to the function $f(t) := t^i$, with respect to the norm

$$\|f\| := \langle f, f \rangle^{1/2}$$

in \mathbf{P}_n . Alternatively, one may construct $(q_i)_0^n$ by its three-term recurrence, an idea popularized specifically for the case of a discrete inner product by Forsythe [2]: One computes

$$\begin{aligned} q_{-1}(t) &= 0, & q_0(t) &= 1, \\ q_i(t) &= (t - \alpha_i)q_{i-1}(t) - \beta_{i-1}^2 q_{i-2}(t), & i &= 1, \dots, n, \end{aligned} \tag{3}$$

with the numbers α_i and β_i computed concurrently by

$$\alpha_i := \langle tq_{i-1}, q_{i-1} \rangle / \langle q_{i-1}, q_{i-1} \rangle, \quad i = 1, \dots, n, \tag{4a}$$

$$\beta_i := \|q_i\| / \|q_{i-1}\|, \quad i = 1, \dots, n-1. \tag{4b}$$

Here, it is assumed that $\langle tf(t), g(t) \rangle = \langle f(t), tg(t) \rangle$.

The computational process (3)-(4) for the vectors α and β is very stable. We will, therefore, have solved Problem A in a satisfactory manner provided we can construct a suitable inner product for which $q_i = p_i$ for $i = n - 1$ and $i = n$. This we now do.

From a computational point of view, the simplest bilinear form \langle , \rangle which is an inner product in P_n is of the form

$$\langle f, g \rangle := \sum_{i=1}^n f(\xi_i) g(\xi_i) w_i, \quad \text{all } f, g, \tag{5}$$

with $\xi_1 < \dots < \xi_n$, and $w_i > 0, i = 1, \dots, n$.

LEMMA 1. Let $(q_i)_0^n$ be the sequence of monic orthogonal polynomials for the inner product (5). Then

$$\prod_{j=1}^n (t - \xi_j) = q_n(t) \tag{6}$$

and

$$w_i = \frac{\gamma}{q_{n-1}(\xi_i) q_n'(\xi_i)}, \quad i = 1, \dots, n, \quad \text{with } \gamma := \frac{\|q_{n-1}\|^2}{\sum_{j=1}^n \frac{q_{n-1}(\xi_j)}{q_n'(\xi_j)}}. \tag{7}$$

Consequently, we can recover (5) from q_{n-1} and q_n .

Proof. The polynomial $q(t) := \prod_{j=1}^n (t - \xi_j)$ is a monic polynomial of degree n which is orthogonal with respect to the inner product (5) to all functions, and hence must agree with q_n . This proves (6). As to (7), we know that q_{n-1} is orthogonal to P_{n-1} . This means that the linear functional L given by the rule

$$Lf := \langle f, q_{n-1} \rangle = \sum_{i=1}^n f(\xi_i) q_{n-1}(\xi_i) w_i, \quad \text{all } f,$$

vanishes on P_{n-1} . Since any $n - 1$ distinct point evaluations are linearly

independent on P_{n-1} , this implies that

$$q_{n-1}(\xi_i) \neq 0, \quad \text{all } i.$$

For the same reason, there is, up to multiplication by a scalar, exactly one linear functional M of the form $Mf = \sum_1^n f(\xi_i)m_i$, all f , which vanishes on P_{n-1} . Since both L and the $(n-1)$ st divided difference $[\xi_1, \dots, \xi_n]$ on the points ξ_1, \dots, ξ_n are such linear functionals, it follows that, for an appropriate scalar γ ,

$$Lf = \gamma[\xi_1, \dots, \xi_n]f = \gamma \sum_{i=1}^n \frac{f(\xi_i)}{\prod_{\substack{j=1 \\ j \neq i}}^n (\xi_i - \xi_j)}, \quad \text{all } f.$$

But this states, with (6), that

$$q_{n-1}(\xi_i)w_i = \gamma/q'_n(\xi_i), \quad i = 1, \dots, n,$$

and, in particular,

$$\|q_{n-1}\|^2 = \gamma \sum_{i=1}^n \frac{q_{n-1}(\xi_i)}{q'_n(\xi_i)},$$

thus proving (7). ■

One may view Lemma 1 as giving a way to construct the computationally simplest inner product with respect to which a given sequence $(p_i)_0^n$ of monic polynomials satisfying a three-term recurrence (2) is orthogonal.

3. A SOLUTION OF PROBLEMS A, B, C

Lemma 1 shows how to reconstruct the discrete inner product (5) from its last two orthogonal polynomials. It also shows the well-known facts that q_n has n real zeros, all simple, and that the $n-1$ zeros of q_{n-1} strictly separate those of q_n . Indeed, the positivity of the w_i 's demands by (7) that $q_{n-1}(\xi_i)q'_n(\xi_i) \text{ signum } \gamma > 0$, all i , while, clearly, $(-)^{n-i}q'_n(\xi_i) \geq 0$, all i , and

therefore

$$q_{n-1}(\xi_i)q_{n-1}(\xi_{i+1}) < 0, \quad i = 1, \dots, n-1,$$

showing q_{n-1} to have a simple zero between any two zeros of q_n .

Conversely, suppose we compute w by

$$w_i := \frac{1}{p_{n-1}(\xi_i) p'_n(\xi_i)}, \quad i = 1, \dots, n, \tag{8a}$$

where

$$\xi_i := \lambda_i, \quad i = 1, \dots, n, \tag{8b}$$

$$p_{n-1}(t) := \prod_{j=1}^{n-1} (t - \mu_j), \tag{8c}$$

$$p_n(t) := \prod_{j=1}^n (t - \lambda_j), \tag{8d}$$

with $\lambda_i < \mu_i < \lambda_{i+1}$, all i . Then $w_i > 0$, all i ; hence (5) is then an inner product on \mathbf{P}_n , and necessarily $p_n = q_n$, by (6), and $p_{n-1} = q_{n-1}$ since $p_{n-1}(\lambda_i) = q_{n-1}(\lambda_i)$, $i = 1, \dots, n$, by (7), and both polynomials are of degree $< n$.

This proves that *Problem A has exactly one solution for given λ and μ satisfying (S)*. Further, since $a = \alpha$ and $b = \beta$ as determined by (3)–(4) depend continuously on ξ and w , while the latter, as determined by (8), depend continuously on λ and μ , it follows that *J depends continuously on λ and μ* .

Problem B is closely related to Problem A. In terms of the monic polynomials

$$p_i(t) = \det(t - J_i), \quad i = 1, \dots, n,$$

we are given the information that

$$\prod_{j=1}^n (t - \lambda_j) = p_n(t) = (t - a_n) p_{n-1}(t) - b_{n-1}^2 p_{n-2}(t)$$

and that

$$\prod_{j=1}^n (t - \lambda_j^*) = p_n^*(t) = (t - a_n^*) p_{n-1}(t) - b_{n-1}^2 p_{n-2}(t).$$

We conclude that

$$\prod_{j=1}^n (t - \lambda_j^*) - \prod_{j=1}^n (t - \lambda_j) = (a_n - a_n^*) p_{n-1}(t),$$

and therefore, comparing coefficients (or else comparing the trace of J with that of J^*),

$$\sum_{j=1}^n (\lambda_j - \lambda_j^*) = a_n - a_n^*.$$

This allows calculation of a_n^* once we know a_n . Further, since we only need to know the weights w for the inner product (5) up to a scalar multiple in order to reconstruct a and b via (3)–(4), it follows that we get J (and uniquely so) by choosing

$$\begin{aligned} \xi_i &:= \lambda_i, \quad i = 1, \dots, n, \\ w_i &:= \frac{-1}{p_n'(\lambda_i) \prod_{j=1}^n (\lambda_i - \lambda_j^*)}. \end{aligned} \quad (9)$$

Note how the assumption $\lambda_i < \lambda_i^* < \lambda_{i+1}$, all i , insures that $w_i > 0$, all i .

The solution of Problem C leads to an intriguing fact which is also of help in the final algorithm for the solution of these problems. We came upon this fact accidentally. We had somehow gained the impression in reading Hochstadt's paper [8] that the correct weights in Lemma 1 would probably be

$$w_i = q_{n-1}(\xi_i) / q_n'(\xi_i), \quad i = 1, \dots, n, \quad (10)$$

and a quick numerical experiment confirmed this guess. Yet, when it came to proving it, we could only prove that $w_i = 1 / [q_{n-1}(\xi_i) q_n'(\xi_i)]$, all i . This seeming contradiction is resolved by consideration of the characteristic polynomials of the *right* principal submatrices of J .

Let S be the permutation matrix carrying $(1, 2, \dots, n)$ into $(n, n-1, \dots, 1)$, i.e.,

$$S := (\delta_{i+j, n+1})_{i,j=1}^n = S^{-1},$$

and denote by \bar{J} the reflection of J across its second diagonal,

$$\bar{J} := S^{-1}JS =: \begin{bmatrix} \bar{a}_1 & \bar{b}_1 & & & & & \\ \bar{b}_1 & \bar{a}_2 & \bar{b}_2 & & & & \\ & & \ddots & \ddots & \ddots & & \\ & & & \bar{b}_{n-2} & \bar{a}_{n-1} & \bar{b}_{n-1} & \\ & & & & \bar{b}_{n-1} & \bar{a}_n & \end{bmatrix}$$

with $\bar{a}_i = a_{n+1-i}$, $\bar{b}_i = b_{n-i}$, all i . Correspondingly, let

$$\bar{p}_{-1}(t) := 0, \quad \bar{p}_0(t) := 1, \quad \bar{p}_i(t) := \det(t - \bar{J}_i), \quad i = 1, \dots, n.$$

LEMMA 2. For $i = 1, \dots, n$, $p_{n-1}(\lambda_i) \bar{p}_{n-1}(\lambda_i) = (b_1 \cdots b_{n-1})^2$.

Proof. For each i , $p_{n-1}(\lambda_i) \bar{p}_{n-1}(\lambda_i)$ is the product of the $(n-1)$ st-order left principal minor with the $(n-1)$ st-order right principal minor of the singular matrix

$$A := \lambda_i - J,$$

i.e.,

$$p_{n-1}(\lambda_i) = \det A \begin{pmatrix} 1, \dots, n-1 \\ 1, \dots, n-1 \end{pmatrix} \quad \text{and} \quad \bar{p}_{n-1}(\lambda_i) = \det A \begin{pmatrix} 2, \dots, n \\ 2, \dots, n \end{pmatrix},$$

and $\det A = 0$. Apply Sylvester's identity (see, e.g., [3, p. 15]), using

$A \begin{pmatrix} 2, \dots, n-1 \\ 2, \dots, n-1 \end{pmatrix}$ as pivotal block, to get

$$0 = \det A \begin{pmatrix} 2, \dots, n-1 \\ 2, \dots, n-1 \end{pmatrix} \det A = \det \begin{bmatrix} \det A \begin{pmatrix} 1, \dots, n-1 \\ 1, \dots, n-1 \end{pmatrix} & \det A \begin{pmatrix} 1, \dots, n-1 \\ 2, \dots, n \end{pmatrix} \\ \det A \begin{pmatrix} 2, \dots, n \\ 1, \dots, n-1 \end{pmatrix} & \det A \begin{pmatrix} 2, \dots, n \\ 2, \dots, n \end{pmatrix} \end{bmatrix}$$

$$= p_{n-1}(\lambda_i) \bar{p}_{n-1}(\lambda_i) - [(-)^{n-1} b_1 \cdots b_{n-1}]^2. \quad \blacksquare$$

Here, we have used the abbreviation

$$A \begin{pmatrix} i_1, \dots, i_r \\ j_1, \dots, j_s \end{pmatrix} := (a_{i_\rho j_\sigma})_{\rho=1, \sigma=1}^{r, s}.$$

If now J is persymmetric, then $J = \bar{J}$ and so $p_i = \bar{p}_i$, all i . The lemma then implies that $[p_{n-1}(\lambda_i)]^2 = (b_1 \cdots b_{n-1})^2$, all i . Since we only need to know the weight vector w up to a scalar multiple, it follows that we only need to know p_n in order to reconstruct a persymmetric J , thus solving Problem C.

We conclude further that the computations (3)–(4) always generate the diagonals α and β of a persymmetric Jacobi matrix if we use the discrete inner product

$$\langle f, g \rangle := \sum_{i=1}^n \frac{f(\xi_i) g(\xi_i)}{\prod_{\substack{j=1 \\ j \neq i}}^n |\xi_i - \xi_j|}.$$

We were interested in Lemma 2 because of its importance for the algorithm in the next section and have therefore not followed the more customary treatment of Problem C. This treatment goes back to Gantmacher and Krein and consists in using the persymmetry of J to construct an equivalent problem of the form B and of half the size, thus reducing it to a problem with a known solution.

4. AN ALGORITHM

Lemma 2 shows that we could also determine the correct weights w for the generation of a and b via (4) by

$$w_i = \bar{p}_{n-1}(\lambda_i) / p'_n(\lambda_i), \quad i = 1, \dots, n.$$

To say it differently, if the inner product (5) is given by

$$\begin{aligned} \xi_i &:= \lambda_i, & i &= 1, \dots, n, \\ w_i &:= p_{n-1}(\lambda_i) / p'_n(\lambda_i), & i &= 1, \dots, n, \end{aligned} \tag{11}$$

then the quantities generated by (3)–(4) are $q_i = \bar{p}_i$, $\alpha_i = \bar{a}_i$, $\beta_i = \bar{b}_i$, all i . This says that use of the weights (11) rather than the weights (8a) in the computations (3)–(4) also generates the nonzero entries of J , but in reverse order. This explains the success in our numerical experiments using the weights (10) referred to earlier: all examples happened to have been persymmetric.

Use of the weights (11) in preference to (8a) has some computational advantages. Because of the interlacing conditions (S), we get the bounds

$$\frac{\lambda_i - \mu_{i-1}}{\lambda_i - \lambda_1} \frac{\mu_i - \lambda_i}{\lambda_n - \lambda_i} < \frac{p_{n-1}(\lambda_i)}{p'_n(\lambda_i)} < 1, \tag{12}$$

where the first (last) factor in the lower bound is to be omitted in case $i = 1$ ($i = n$). This shows that overflow or underflow is highly unlikely to occur in the calculation of the weights (11). By contrast, the computation of the numbers $1 / [p_{n-1}(\lambda_i) p'_n(\lambda_i)]$ has to be carefully monitored, in general, for the occurrence of overflow or underflow, else one has to compute the logarithms of these numbers, a somewhat more expensive procedure.

We offer the following algorithm for the solution of Problem A, and recall that Problems B and C can also be solved by it, if the definition of $p_{n-1}(\lambda_i) := \prod_1^{n-1} (\lambda_i - \mu_j)$ used here is modified appropriately.

ALGORITHM. Given the n eigenvalues $\lambda_1 < \dots < \lambda_n$ of the Jacobi matrix (1) and the $n - 1$ eigenvalues $\mu_1 < \dots < \mu_{n-1}$ of its left principal minor of order $n - 1$. Note that, necessarily, $\lambda_i < \mu_i < \lambda_{i+1}$, all i .

1. Compute the weights w from λ and μ :
 - 1.1 $\text{temp}(i-1) := \lambda_i, i = 2, \dots, n$
 - 1.2 for $i = 1, \dots, n$, do:
 - 1.21 $w_i := \prod_{j=1}^{n-1} (\lambda_i - \mu_j) / [\lambda_i - \text{temp}(j)]$
 - 1.22 $\text{temp}(i) := \lambda_i$
2. Generate the values at λ of the first two orthogonal polynomials:
 - 2.1 $s := \sum_{j=1}^n w_j = \langle \bar{p}_0, \bar{p}_0 \rangle$
 - 2.2 $\bar{a}_1 := (\sum_{j=1}^n w_j \lambda_j) / s = \langle \bar{p}_0, \bar{p}_1 \rangle / s$
 - 2.3 for $i = 1, \dots, n$, do:
 - 2.31 $\text{pkml}(i) := 1 = \bar{p}_0(\lambda_i)$
 - 2.32 $\text{pk}(i) := \lambda_i - \bar{a}_1 = \bar{p}_1(\lambda_i)$
3. Compute $\|\bar{p}_k\|^2$ and $\bar{a}_k, \bar{b}_{k-1}^2$, then use them to generate the values at λ of \bar{p}_{k+1} from those of \bar{p}_k and \bar{p}_{k-1} by the three-term recurrence:
 - 3.1 for $k = 2, \dots, n$, do:
 - 3.11 $s' := s = \|\bar{p}_{k-1}\|^2$
 - 3.12 $s := t := 0$
 - 3.13 for $i = 1, \dots, n$, do:
 - 3.131 $p := w_i \times \text{pk}(i)^2$
 - 3.132 $s := s + p$
 - 3.133 $t := t + \lambda_i p$
 - 3.14 $\bar{b}_{k-1}^2 := s / s'$
 - 3.15 $\bar{a}_k := t / s$
 - 3.16 for $i = 1, \dots, n$, do:
 - 3.161 $p := \text{pk}(i)$
 - 3.162 $\text{pk}(i) := (\lambda_i - \bar{a}_k) p - \bar{b}_{k-1}^2 \text{pkml}(i)$
 - 3.163 $\text{pkml}(i) := p$
4. Compute \bar{b}_k from \bar{b}_k^2 . Also, if a and b , rather than the vectors \bar{a} and \bar{b} , are wanted, this is the place to reorder them.
 - 4.1 $\bar{b}_k := \text{sqrt}(\bar{b}_k^2), k = 1, \dots, n-1$

Output consists of the vectors \bar{a} and \bar{b} , with $\bar{a}_i = a_{n+1-i}, \bar{b}_i = b_{n-i}$, all i , and a and b the diagonals of (1).

We have carried out various numerical experiments with this algorithm and describe here only three.

For the n th-order Jacobi matrix J_n with general row

$$1, \quad -2, \quad 1, \quad \quad \quad (i)$$

the eigenvalues are given explicitly (and in order) by the formula

$$\lambda_j = 2 \left(\cos \frac{j\pi}{n+1} - 1 \right), \quad j = 1, \dots, n.$$

Starting with these values and the corresponding sequence $(\mu_j)_1^{n-1}$ for J_{n-1} , the algorithm produced approximations to the diagonal and the off-diagonal entries of J_n whose maximum and average error are recorded for $n = 25, 50, 100,$ and 200 in the first columns of the Table 1. All calculations were carried out on a UNIVAC 1110 in single precision (27-bit floating-point mantissa).

For variety, we also consider the n -th order Jacobi matrix J_n with general row

$$1 - \frac{i}{n}, \quad \frac{i}{n} - 2, \quad 1 - \frac{i+1}{n}, \quad i = 1, \dots, n. \tag{ii}$$

We know of no simple formula for its eigenvalues, and therefore used the algorithm tq1 1 on pp. 232–233 of Wilkinson and Reinsch’s handbook [10] to compute them and those of J_{n-1} . The tolerance (relative error requirement) for tq1 1 we chose as $1. - 7$. With this spectral information, we entered the above algorithm and so reconstructed J_n approximately. Errors of this reconstruction are also given in Table 1, in the last four columns. There is a significant deterioration as n increases.

As can be expected from the formula (11) for the weights (w_i) , the condition of the problem of determining J_n from (λ_i) and (μ_i) deteriorates as one or more μ_i approach the corresponding λ_i , since then one or more of the weights approach zero. This is shown even more strikingly when the matrix of the last example is reflected across its second diagonal, i.e., when the

TABLE 1
 MAXIMUM AND AVERAGE ERROR IN THE DIAGONAL AND OFF-DIAGONAL ENTRIES OF TWO SPECIFIC JACOBI MATRICES^a

<i>n</i>	(i)				(ii)			
	Diagonal		Off-Diagonal		Diagonal		Off-Diagonal	
	Max.	Ave.	Max.	Ave.	Max.	Ave.	Max.	Ave.
25	4. - 7	2. - 7	2. - 7	6. - 8	2. - 6	5. - 7	1. - 6	4. - 7
50	9. - 7	1. - 7	4. - 7	2. - 7	6. - 4	1. - 5	5. - 5	2. - 6
100	2. - 6	7. - 7	8. - 7	2. - 7	7. - 1	3. - 2	3. - 1	1. - 2
200	3. - 6	9. - 7	1. - 6	3. - 7	8. - 1	2. - 2	5. - 1	1. - 2

^aAs reconstructed with the algorithm of this section from (approximate) spectral data.

Jacobi matrix with the general row

$$1 - (n + 1 - i)/n, \quad (n + 1 - i)/n - 2, \quad 1 - (n - i)/n, \quad i = 1, \dots, n,$$

is considered. Now the reconstruction breaks down in single precision already for $n = 30$, since $\mu_1 - \lambda_1$ becomes too small. Even for $n = 20$, we have $\mu_1 - \lambda_1 \sim 2 \cdot 10^{-7}$. In fact, in computations using tql 1 to obtain the spectral information, some weights become negative for $n = 30$, while for $n = 10$ and 20 we obtain approximations with errors of the order of $1 \cdot 10^{-4}$ and $2 \cdot 10^{-2}$, respectively.

5. THE CONNECTION WITH GAUSS QUADRATURE

For the given Jacobi matrix J in (1), let $(P_i)_0^n$ be the polynomial sequence generated by the recursion

$$\begin{aligned} b_{i+1}P_{i+1}(t) &= (t - a_{i+1})P_i(t) - b_iP_{i-1}(t), \quad i = 0, \dots, n - 1 \\ P_{-1}(t) &:= 0, \quad P_0(t) := 1, \end{aligned} \tag{13}$$

with b_0 arbitrary, $b_n \neq 0$. The sequence (P_i) is related to the sequence (p_i) with $p_i(t) := \det(t - J_i)$, all i , of monic polynomials by

$$p_i = (b_1 \cdots b_i)P_i, \tag{14}$$

as one verifies easily, e.g., by comparing (13) and (2).

Let now ω be a monotone function on some interval $[A, B]$ so that $(P_i)_0^n$ is *orthonormal* with respect to the inner product

$$\langle f, g \rangle_\omega := \int_A^B f(t)g(t)\omega(dt).$$

(Lemma 1 provides a simple proof of the existence of such ω .) Then the zeros $\lambda_1 < \dots < \lambda_n$ of P_n must lie in $[A, B]$, and there exist positive weights w_1, \dots, w_n such that for every $f \in C^{2n}[A, B]$,

$$\int_A^B f(t)\omega(dt) = \sum_{j=1}^n w_j f(\lambda_j) + \frac{b_1 \cdots b_n f^{(2n)}(\xi)}{(2n)!} \quad \text{for some } \xi \in]A, B[. \tag{15}$$

If we take this fact for granted, then it follows that

$$\delta_{i-j} = \langle P_i, P_j \rangle_\omega = \sum_{r=1}^n w_r P_i(\lambda_r) P_j(\lambda_r) \quad \text{for } i+j < 2n,$$

showing that

$$\langle f, g \rangle := \sum_{i=1}^n f(\lambda_i) g(\lambda_i) w_i \tag{5}$$

is an inner product on P_n with respect to which $(P_i)_0^{n-1}$ is orthonormal, and hence for which $(p_i)_0^n$ is an orthogonal sequence.

This shows that the construction of the weights (w_i) for (5), which was crucial for our numerical solution of the various inverse eigenvalue problems, can be started from any convenient formula for the weights in a Gaussian quadrature formula.

For instance, one could start with the following consequence of the Christoffel-Darboux formula:

$$w_j P^T(\lambda_j) P(\lambda_j) = 1 \quad \text{for } j = 1, \dots, n. \tag{16}$$

Here, $P(\lambda)$ denotes the n -vector $(P_0(\lambda), \dots, P_{n-1}(\lambda))$. By (13), $P(\lambda_j)$ is an eigenvector for J belonging to the eigenvalue λ_j . Therefore, with $u_j := (u_{1j}, \dots, u_{nj})$ a unit eigenvector of J for λ_j , (16) implies that

$$w_j = u_{ij}^2 / P_{i-1}^2(\lambda_j), \quad i = 1, \dots, n. \tag{17}$$

Since $P_0(\lambda) = 1$, we obtain in this way the formula

$$w_j = u_{1j}^2, \quad j = 1, \dots, n, \tag{18}$$

used by Golub and Welsch [4] to compute the weights. Problems A, B, and C can now be solved by deriving from the given data information about the eigenvectors of J .

A more direct approach might be to start with the well-known formula

$$w_j = -\frac{k_{n+1}}{k_n} \frac{1}{P_{n+1}(\lambda_j) P'_n(\lambda_j)}, \quad j = 1, \dots, n, \tag{19}$$

with k_j the leading coefficient of P_j , i.e., $k_j = 1/(b_1 \cdots b_j)$. This formula involves the “next” orthogonal polynomial P_{n+1} . But, since $P_n(\lambda_j) = 0$ for all j ,

we have

$$P_{n+1}(\lambda_j)/k_{n+1} = p_{n+1}(\lambda_j) = -b_n^2 p_{n-1}(\lambda_j) = -b_n^2 P_{n-1}(\lambda_j)/k_{n-1}$$

by the three-term recurrence; therefore we also have

$$w_j = \frac{1}{b_n P_{n-1}(\lambda_j) P_n'(\lambda_j)}, \quad j=1, \dots, n, \quad (19')$$

which shows how (7) could have been derived from standard results in Gauss quadrature.

The authors are very pleased to acknowledge stimulating discussions with Professor Victor Barcilon of the University of Chicago.

The authors wish to thank Mr. Daniel Boley of Stanford University for corroborating some of their calculations.

The work of G. H. G. was begun when he was a guest worker at the Mathematics Research Center, whose hospitality he greatly enjoyed.

REFERENCES

- 1 V. Barcilon, Iterative solutions of inverse eigenvalue problems for Jacobi matrices, unpublished manuscript.
- 2 George E. Forsythe, Generation and use of orthogonal polynomials for data-fitting with a digital computer, *J. SIAM* 5 (1957) 74–88.
- 3 F. R. Gantmacher and M. G. Krein, *Oszillationsmatrizen, Oszillationskerne und kleine Schwingungen mechanischer Systeme*, Akademie-Verlag, Berlin, 1960.
- 4 G. H. Golub and J. H. Welsch, Calculation of Gauss quadrature rules, *Math. Comp.* 23 (1969) 221–230.
- 5 L. J. Gray and D. G. Wilson, Construction of a Jacobi matrix from spectral data, *Linear Algebra Appl.* 14 (1976) 131–134.
- 6 Ole H. Hald, Inverse eigenvalue problems for Jacobi matrices, *Linear Algebra Appl.* 14 (1976) 63–85.
- 7 Harry Hochstadt, On some inverse problems in matrix theory, *Arch. Math.* 18 (1967) 201–207.
- 8 Harry Hochstadt, On the construction of a Jacobi matrix from spectral data, *Linear Algebra Appl.* 8 (1974) 435–446.
- 9 Burton Wendroff, On orthogonal polynomials, *Proc. Amer. Math. Soc.* 12 (1961) 554–555.
- 10 J. H. Wilkinson and C. Reinsch, *Linear Algebra, Handbook for Automatic Computation II*, Springer, Berlin, 1971.

Received 25 February 1977; revised 12 April 1977