

COMPLEMENTARITY AND RELATED PROBLEMS: A SURVEY

Michael C. Ferris^{1,2} and Christian Kanzow^{3,4}

¹ University of Wisconsin – Madison
Computer Sciences Department
1210 West Dayton Street
Madison, WI 53706
USA
e-mail: ferris@cs.wisc.edu

³ University of Hamburg
Institute of Applied Mathematics
Bundesstrasse 55
20146 Hamburg
Germany
e-mail: kanzow@math.uni-hamburg.de

November 23, 1998

Abstract. This survey gives an introduction to some of the recent developments in the field of complementarity and related problems. After presenting two typical examples and the basic existence and uniqueness results, we focus on some new trends for solving nonlinear complementarity problems. Extensions to mixed complementarity problems, variational inequalities and mathematical programs with equilibrium constraints are also discussed.

Key Words. Nonlinear complementarity problem; Mixed complementarity problem; Variational inequalities; Mathematical programs with equilibrium constraints.

²The research of this author was partially supported by National Science Foundation Grants CCR-9619765 and CDA-9726385.

⁴Current address (October 1, 1998 – March 31, 1999): Dipartimento di Informatica e Sistemistica, Università di Roma “La Sapienza”, Via Buonarroti 12, 00185 Roma, Italy; e-mail: ckanzow@dis.uniroma1.it. The research of this author was supported by the DFG (Deutsche Forschungsgemeinschaft).

1 Introduction

In this paper, we give a survey of complementarity problems that covers some standard results and presents a number of new developments in the field. These developments are first described for the most basic problem, the nonlinear complementarity problem (NCP), which is to find a vector $x \in \mathbb{R}^n$ satisfying the system of equations and inequalities

$$x \geq 0, \quad F(x) \geq 0, \quad x^T F(x) = 0$$

or, equivalently,

$$x_i \geq 0, \quad F_i(x) \geq 0, \quad x_i F_i(x) = 0 \quad \text{for } i = 1, \dots, n.$$

Here $F : X \rightarrow \mathbb{R}^n$ is a given function defined on a subset $X \subseteq \mathbb{R}^n$ containing at least the nonnegative orthant.

Due to expository and space considerations, we do not always present a result in its most general form; for example, we will introduce the class of monotone functions in Section 3 and state a couple of important results for this class even though some of them actually hold for the somewhat broader classes of pseudomonotone or P_0 -functions.

Many of the basic theoretical results in complementarity problems have been known for a long time; an excellent survey of much of the research in this area prior to 1990 can be found in [39]. Further references and more recent work can also be found in [31, 38, 75]. Here, we give just a flavor of the known existence and uniqueness results and concentrate more on the algorithmic point of view.

The organization of this paper is as follows. Section 2 gives two typical applications of complementarity problems as they arise in economics and engineering sciences. In Section 3 we then state a few existence and uniqueness results for the standard NCP, and Section 4 describes a couple of different solution techniques for NCP. Section 5 deals with some more general equilibrium-type problems, namely mixed complementarity problems, variational inequalities and mathematical programs with equilibrium constraints.

The notation used in this paper is assumed to be rather standard: Then n -dimensional Euclidean space is denoted by \mathbb{R}^n with $\|x\|$ being the Euclidean norm of a vector $x \in \mathbb{R}^n$. The scalar product of two vectors $x, y \in \mathbb{R}^n$ is written as $x^T y$. An inequality $x \geq 0$ means that all components of the vector $x \in \mathbb{R}^n$ are nonnegative. If $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a continuously differentiable function, we denote its Jacobian at a point $x \in \mathbb{R}^n$ by $F'(x)$. Furthermore, if $M \in \mathbb{R}^{n \times n}$ denotes an arbitrary matrix with elements m_{ij} and $I, J \subseteq \{1, \dots, n\}$ are two given subsets, then M_{IJ} is an $|I| \times |J|$ -dimensional submatrix with entries m_{ij} for $i \in I$ and $j \in J$.

2 Examples

Complementarity problems arise frequently in the general equilibrium theory [2] of economics. Many complex computable general equilibrium models are used for various aspects of policy design and analysis, including carbon abatement [60] and trade reform [40]. Other

economic applications use game theory [68]; new examples of this are becoming popular due to deregularization of electricity markets [92].

Our particular example is taken from [88] and is a schematic Arrow-Debreu [2] equilibrium problem. The economy contains ℓ commodities, m utility-maximizing consumers and n profit-maximizing producers. The key variables of the model are the prices p of the commodities. Given these prices, it is assumed that the agents demand certain amounts of the commodities, which are supplied so as to maximize profit in each production sector. Complementarity is derived from Walras' law, namely that supply will equal demand in the equilibrium state.

The production side of the economy depends on a technology matrix

$$A = [a^1 \quad a^2 \quad \dots \quad a^n] \in \mathbb{R}^{\ell \times n}$$

that determines the unit output level of each sector. Generally, the entries of A are nonlinear functions of the price vector p and can be calculated using production functions (see [32]). For simplicity, we will assume here a constant technology matrix, and convert the activity level z_j of sector j into commodities using $a^j z_j$. Under the aforementioned assumption of price taking, each sector maximizes profits for given prices p :

$$\max\{p^T a^j z_j : z_j \geq 0\}.$$

The necessary and sufficient optimality conditions for production are therefore:

$$z \geq 0, \quad A^T p \leq 0, \quad p^T A z = 0. \quad (1)$$

For consumption, we derive a demand function for each of the consumers in the economy. The i th consumer is endowed with an initial wealth w_i and, assuming that he is also a price taker, will maximize his utility subject to his budget constraints. As is typical in such problems, the utility function is assumed to be either Cobb-Douglas

$$u^i(d) := \prod_c d_c^{\lambda_c}, \quad \lambda_c \geq 0, \quad \sum_c \lambda_c = 1,$$

or constant elasticity of substitution (CES)

$$u^i(d) := \left(\sum_{c=1}^{\ell} \lambda_c x_c^\rho \right)^{1/\rho}, \quad \lambda_c \geq 0, \quad c = 1, \dots, \ell,$$

where λ_c are given parameters dependent on the consumer i . The consumer solves

$$\max\{u^i(d^i) : p^T d^i \leq w_i, \quad d^i \geq 0\}, \quad (2)$$

which in the case of Cobb-Douglas utility, gives

$$d_c^i(p, w_i) = \frac{\lambda_c w_i}{p_c}, \quad (3)$$

whereas for CES utility,

$$d_c^i(p, w_i) = \frac{(p_c/\lambda_c)^{r-1} w_i}{\sum_{j=1}^{\ell} p_j^r/\lambda_j^{r-1}}, \quad r = \rho/(\rho - 1). \quad (4)$$

Typically, wealth is determined from an initial endowment of commodities e^i , so that $w_i = p^T e^i$.

While the prices were assumed above to be given to each of the agents in the economy, the model actually uses Walras' law to determine the prices so that markets clear. This clearance condition is stated succinctly as follows:

$$p \geq 0, \quad Az - \sum_{i=1}^m (d^i(p, p^T e^i) - e^i) \geq 0, \quad p^T (Az - \sum_{i=1}^m (d^i(p, p^T e^i) - e^i)) = 0.$$

If the price p is positive then supply $Az + \sum_{i=1}^m e^i$ must equal demand $\sum_{i=1}^m d^i(p, w_i)$. Otherwise, when excess supply is available, the price must be zero.

Summing all the information regarding production and consumption results in the following complementarity problem:

$$z \geq 0, \quad -A^T p \geq 0, \quad p^T Az = 0, \\ p \geq 0, \quad Az - \sum_{i=1}^m (d^i(p, p^T e^i) - e^i) \geq 0, \quad p^T (Az - \sum_{i=1}^m (d^i(p, p^T e^i) - e^i)) = 0.$$

Note that if (p, z) solves the above problem, then so does $(\lambda p, z)$ for any $\lambda > 0$. This frequently causes algorithms difficulty in determining a solution. Typically this singularity is removed by fixing a numeraire, that is fixing a price and dropping the corresponding complementary relationship.

The data for a particular instance of this model is available in MCPLIB [20]. Many extensions of this basic framework are possible [32, 86, 87]; examples using specialized software are described in [27, 29, 86, 87].

Another example of the use of complementarity problems is in the field of structural analysis. One of the aims of structural analysis is to find displacement, stress and strain fields in a given structure under given conditions. Typically, loads F are specified and the problem is to find displacements u , stresses Q and strains q . Compatibility or kinematic relationships relate u and q , equilibrium or statics relationships relate F and Q and the constitutive laws relate Q and q . There are many models in use by engineers due to different assumptions for displacement and constitutive laws. For example, small or large displacements can be assumed, different discretization or finite element models can be applied and time independent plasticity relationships can be used or not.

Starting in the 1970's, Maier and colleagues [54, 55, 56, 99, 100] proposed the use of complementarity in mathematical models for structural mechanics. The general form of the model is as follows. The compatibility relationships are determined from the geometry of the structure using $q = q(u)$. The equilibrium relationships are normally given by

$$C^T Q = F, \quad C = \frac{\partial q}{\partial u},$$

so that in a small displacement model $q = Cu$. The constitutive laws can be written

$$Q = Se, \quad p = N\lambda, \quad q = e + p,$$

where e and p represent elastic and plastic strains, S corresponds to elastic stiffnesses for each member of the structure, N contains unit outward normals and λ are the plastic multipliers. The yield function satisfies the following complementarity relationship

$$\lambda \geq 0, \quad -N^T Q + Q_* + h(\lambda) \geq 0, \quad \lambda^T (-N^T Q + Q_* + h(\lambda)) = 0.$$

The elastic limit Q_* , and h (representing softening behavior) are material properties and are typically given. Most of the previous work in this area has been concerned with linear models of this format; typically, many of the above variables are eliminated to construct a standard form linear complementarity model. A more reasonable modeling approach maintains the sparsity and the mixed complementarity nature of the above formulation (see Section 5.1 and [98]). Nonlinear extensions of the models are detailed in [7, 32, 100].

3 Existence

In this section, we deal with the following questions: Does the NCP have at least one solution? When is the solution unique? What is the structure of the solution set?

In general, it is impossible to give a satisfactory answer to any of these questions without posing certain conditions on the function F . In order to give an idea of what can happen, consider the following simple examples.

Example 3.1 *In all examples given here, the function $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is defined for $n = 1$.*

- (a) *If $F(x) = -x - 1$, then NCP has no solution since $F(x) < 0$ for all $x \geq 0$.*
- (b) *If $F(x) = x - 1$, then NCP has a unique solution $x^* = 1$.*
- (c) *If $F(x) \equiv 0$, then NCP has infinitely many solutions, namely the half line $[0, +\infty)$.*
- (d) *If $F(x) = (x-1)^2 - 1$, then NCP has two isolated solutions, namely $x^* = 0$ and $x^{**} = 2$.*

In order to state some existence and, possibly, uniqueness results, we now introduce some classes of functions which play the role of corresponding convexity properties in optimization problems.

Definition 3.2 *Let $X \subseteq \mathbb{R}^n$ be a nonempty set and $F : X \rightarrow \mathbb{R}^n$ be a given function. Then F is said to be*

- (i) *monotone (on X) if $(x - y)^T (F(x) - F(y)) \geq 0$ for all $x, y \in X$;*
- (ii) *strictly monotone (on X) if $(x - y)^T (F(x) - F(y)) > 0$ for all $x, y \in X$ with $x \neq y$;*
- (iii) *strongly monotone (on X) if there exists a constant $\mu > 0$ (called modulus) such that $(x - y)^T (F(x) - F(y)) \geq \mu \|x - y\|^2$ for all $x, y \in X$.*

Using these definitions, we can state the following result.

Theorem 3.3 *Let $F : X \rightarrow \mathbb{R}^n$ be a continuous function on $X = \mathbb{R}_+^n$. Then the following statements hold:*

- (a) *If F is monotone on X , then NCP has a convex (possibly empty) solution set.*
- (b) *If F is strictly monotone on X , then NCP has at most one solution.*
- (c) *If F is strongly monotone on X , then NCP has a unique solution.*

Unfortunately, the assumption in Theorem 3.3 (c) is rather strong and usually not satisfied in many practical applications. On the other hand, the monotonicity assumption in Theorem 3.3 (a) is a much weaker condition which, in fact, can be guaranteed to hold in quite a few situations. So it would be desirable to have an existence result also for monotone complementarity problems. Fortunately, such a result exists if, in addition, we assume that there is a strictly feasible point.

Definition 3.4 *A vector $\hat{x} \in \mathbb{R}^n$ is called strictly feasible for NCP if $\hat{x} > 0$ and $F(\hat{x}) > 0$.*

The strict feasibility assumption is closely related to the well-known Slater condition in convex optimization problems and allows us to state the following result.

Theorem 3.5 *Let $F : X \rightarrow \mathbb{R}^n$ be a continuous function on $X = \mathbb{R}_+^n$. Assume that F is monotone on X and there is a strictly feasible point \hat{x} for NCP. Then NCP has a nonempty and compact solution set.*

Proofs of most of the results stated in this section can be found in [66], see also [39, 17] for some generalizations and specializations.

4 Algorithms

Here we want to describe the basic ideas of some solution methods for the nonlinear complementarity problem. Suitable algorithms should have the following desirable properties:

1. globally convergent
2. locally fast convergent.

The first requirement says that, if possible, we want to have a method which is guaranteed to find a solution of the NCP independent of how good or bad our starting point is. This can usually be guaranteed only if F has certain properties like being monotone or strongly monotone.

For the second point, a method is viewed as being locally fast convergent if it is locally superlinearly or quadratically convergent. Fast convergence can often be established under a condition called strong regularity [81].

Definition 4.1 Let x^* be a solution of NCP and define the index sets

$$\alpha := \{i \mid x_i^* > 0 = F_i(x^*)\} \quad \text{and} \quad \beta := \{i \mid x_i^* = 0 = F_i(x^*)\}.$$

Then x^* is called a strongly regular solution if the submatrix $F'(x^*)_{\alpha\alpha}$ is nonsingular and the Schur-complement

$$F'(x^*)_{\alpha\cup\beta, \alpha\cup\beta} / F'(x^*)_{\alpha\alpha} := F'(x^*)_{\beta\beta} - F'(x^*)_{\beta\alpha} F'(x^*)_{\alpha\alpha}^{-1} F'(x^*)_{\alpha\beta} \in \mathbb{R}^{|\beta| \times |\beta|}$$

has positive principal minors.

The next subsections discuss several different algorithms. Most of these algorithms are based on a suitable reformulation of the complementarity problem as a nonsmooth equation.

4.1 Projection-Type Methods

Projection-type methods are based on the following simple observation.

Proposition 4.2 A vector x^* is a solution of NCP if and only if x^* satisfies the fixed-point equation

$$x = (x - \gamma F(x))_+,$$

where $\gamma > 0$ is an arbitrary constant and z_+ denotes the Euclidean projection of a vector $z \in \mathbb{R}^n$ onto the nonnegative orthant.

Projection-type methods typically attempt a fixed-point iteration. The simplest consists of the iteration

$$x^{k+1} := (x^k - \gamma F(x^k))_+, \quad k = 0, 1, 2, \dots, \quad (5)$$

where $x^0 \in \mathbb{R}_+^n$ is any given starting point, see [4, 5, 91]. Using Banach's fixed point theorem, it follows easily that this method is globally convergent if F is strongly monotone and globally Lipschitzian provided the constant $\gamma > 0$ is sufficiently small. Furthermore, the rate of convergence one can expect is linear.

Modifications of this projection method have better global convergence properties. For example, the extragradient method [49, 50, 62, 67] generates iterates using the formula

$$x^{k+1} := (x^k - \gamma F((x^k - \gamma F(x^k))_+))_+, \quad k = 0, 1, 2, \dots, \quad (6)$$

i.e., the difference with (5) is that the function F is not evaluated at the previous iterate x^k , but at a point which would already be the next iterate for (5).

The extragradient method can be shown to be globally convergent if F is monotone (rather than strongly monotone) and globally Lipschitzian provided the constant $\gamma > 0$ is sufficiently small, see [67].

In fact, it is even possible to remove the assumptions that F is globally Lipschitzian and $\gamma > 0$ has to be sufficiently small (in practice, it is usually not known how small γ has to be). This is usually done by choosing γ dynamically in a suitable line search within the extragradient method, see [94, 95, 41, 93].

These modified extragradient methods have nice global convergence properties for monotone complementarity problems, but they are at best linearly convergent. Due to their slow convergence, we recommend their use only if either the complementarity problem is huge or if it is impractical to form F' .

4.2 Merit Functions

We first give the definition of a merit function as it will be used within this survey.

Definition 4.3 *A function $\Psi : \mathbb{R}^n \rightarrow \mathbb{R}$ is called a merit function for NCP if it has the following properties:*

- (a) $\Psi(x) \geq 0$ for all $x \in \mathbb{R}^n$;
- (b) $\Psi(x) = 0$ if and only if x is a solution of NCP.

This definition suggests replacing the NCP by the unconstrained minimization problem

$$\min \Psi(x), \quad x \in \mathbb{R}^n.$$

In order to apply standard optimization tools to this minimization problem, the merit function should satisfy certain smoothness conditions. In fact, the merit function Ψ should be at least once continuously differentiable, if possible even twice. It turns out, however, that most known merit functions are only once continuously differentiable. For example, the implicit Lagrangian [59]

$$\Psi(x) := \sum_{i=1}^n \left\{ x_i F_i(x) + \frac{1}{2\alpha} (\max^2\{0, x_i - \alpha F_i(x)\} - x_i^2 + \max^2\{0, F_i(x) - \alpha x_i\} - F_i(x)^2) \right\} \quad (7)$$

(where $\alpha > 1$ is a fixed parameter), the Fischer-Burmeister merit function [26, 37]

$$\Psi(x) := \frac{1}{2} \sum_{i=1}^n \left(\sqrt{x_i^2 + F_i(x)^2} - x_i - F_i(x) \right)^2 \quad (8)$$

or the penalized Fischer-Burmeister merit function [10]

$$\Psi(x) := \frac{1}{2} \sum_{i=1}^n \left\{ \lambda \left(\sqrt{x_i^2 + F_i(x)^2} - x_i - F_i(x) \right) - (1 - \lambda) \max\{0, x_i\} \max\{0, F_i(x)\} \right\}^2 \quad (9)$$

(where $\lambda \in (0, 1)$ is a fixed parameter) are all known to be merit functions that are once but not twice continuously differentiable.

This is one of the reasons why we do not suggest minimizing a merit function directly using existing unconstrained optimization software. Another reason is that most merit functions can be derived from corresponding equation reformulations (see the following subsection), and it is the interplay between such an equation reformulation and its merit function that makes these methods successful.

Apart from this, merit functions are often used in order to monitor global convergence of different algorithms. We believe this is actually the major contribution of a merit function.

We close this subsection by stating two types of results for the above mentioned three merit functions. The first result indicates sufficient conditions to ensure that a stationary point of the merit function Ψ is a global minimum and hence a solution of NCP.

Theorem 4.4 *Assume that $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuously differentiable. If F is strongly monotone, then every stationary point of the implicit Lagrangian (7) is a solution of NCP. The same statement holds for the Fischer-Burmeister merit function (8) and the penalized Fischer-Burmeister merit function (9) under the weaker assumption that F is monotone.*

Most algorithms guarantee that any accumulation point of the iterates is at least a stationary point of Ψ and therefore, under the conditions of Theorem 4.4, a solution of the underlying complementarity problem. The next result concerns the compactness of the level sets

$$\mathcal{L}(c) := \{x \in \mathbb{R}^n \mid \Psi(x) \leq c\}$$

for a given constant c . This is a critical result, since a compact level set implies that the sequence of iterates will have an accumulation point.

Theorem 4.5 *Assume that $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and let $c \in \mathbb{R}$ be any given constant. If F is strongly monotone, then the level sets $\mathcal{L}(c)$ for any of the three merit functions (7)–(9) are compact (possibly empty). The same statement holds for the penalized Fischer-Burmeister merit function (9) under the weaker assumption that F is monotone and NCP has a strictly feasible point.*

There have been many other merit functions proposed in the literature, see e.g. [53, 48, 46, 96]. Currently, the penalized Fischer-Burmeister merit function and those in [48, 96] have the strongest properties.

4.3 Equation-Based Methods

It is possible to reformulate the NCP as a nonlinear system of equations using so-called NCP-functions.

Definition 4.6 *A function $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ is called an NCP-function if it satisfies the condition*

$$\phi(a, b) = 0 \iff a \geq 0, b \geq 0, ab = 0.$$

The following functions are NCP-functions:

- (a) $\phi(a, b) = \min\{a, b\}$;
- (b) $\phi(a, b) = \sqrt{a^2 + b^2} - a - b$;
- (c) $\phi(a, b) = \lambda (\sqrt{a^2 + b^2} - a - b) - (1 - \lambda) \max\{0, a\} \max\{0, b\}$ with $\lambda \in (0, 1)$ fixed;
- (d) $\phi(a, b) = -ab + \frac{1}{2} \min^2\{0, a + b\}$.

Normally (a) is called the minimum function, and (b) is called the Fischer-Burmeister function (it first appeared in a paper by Fischer [33] where it is attributed to Burmeister). Example (c) is termed the penalized Fischer-Burmeister function [10]. Example (d) is the only NCP-function listed that is differentiable on the whole space \mathbb{R}^2 [24]. Many other NCP-functions are known in the literature, see [57, 96].

For any given NCP-function ϕ , define $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$\Phi(x) := \begin{pmatrix} \phi(x_1, F_1(x)) \\ \vdots \\ \phi(x_n, F_n(x)) \end{pmatrix}. \quad (10)$$

We call Φ the equation operator corresponding to ϕ . As an immediate consequence of the definition of Φ , we obtain the following characterization of the complementarity problem.

Proposition 4.7 *Let ϕ be an NCP-function and Φ be the corresponding equation operator defined by (10). Then x^* is a solution of NCP if and only if x^* is a solution of the nonlinear system of equations $\Phi(x) = 0$.*

If F and the NCP-function ϕ are continuously differentiable, then the same holds for the equation operator Φ . In this case, we can apply the classical Newton method to $\Phi(x) = 0$ to solve the underlying NCP. This leads to the iteration

$$x^{k+1} := x^k - \Phi'(x^k)^{-1} \Phi(x^k), \quad k = 0, 1, 2, \dots$$

for a given starting point $x^0 \in \mathbb{R}^n$. We can expect local fast convergence if the standard assumptions for the Newton method are satisfied; in particular, the Jacobian matrix $\Phi'(x^*)$ has to be nonsingular at a solution x^* of NCP. Unfortunately, the following result from [45] shows that this matrix is singular at any degenerate solution. (Here, a solution x^* is called *degenerate* if the index set β from Definition 4.1 is nonempty.)

Proposition 4.8 *Let x^* be a degenerate solution of NCP. Assume that F and the NCP-function ϕ are differentiable, and let Φ denote the corresponding equation operator. Then the Jacobian $\Phi'(x^*)$ is singular.*

One way to overcome the singularity problem is to use a nondifferentiable NCP-function. This, in general, leads to a nondifferentiable equation operator Φ . However, Φ is usually at least locally Lipschitzian, so Φ is almost everywhere differentiable by a result due to Rademacher. Let us denote by D_Φ the set of differentiable points of Φ . Then we define the so-called *B-subdifferential* [82]

$$\partial_B \Phi(x) := \{H \in \mathbb{R}^{n \times n} \mid \exists \{x^k\} \subseteq D_\Phi : x^k \rightarrow x \text{ and } \Phi'(x^k) \rightarrow H\}$$

as well as its convex hull

$$\partial \Phi(x) := \text{conv} \{ \partial_B \Phi(x) \}$$

which is called the *generalized Jacobian* [15]. Elements from $\partial_B \Phi(x)$ or $\partial \Phi(x)$ can be used instead of the classical (and in our situation sometimes not existing) Jacobian $\Phi'(x)$, and a typical nonsmooth Newton method then leads to the iteration

$$x^{k+1} := x^k - H_k^{-1} \Phi(x^k), \quad k = 0, 1, 2, \dots, \quad (11)$$

where H_k is an arbitrary element from $\partial_B \Phi(x^k)$.

Although, in general, it is difficult to compute elements from the B-subdifferential, we are interested only in particular operators Φ , and for those it is usually not difficult to compute an element $H_k \in \partial_B \Phi(x)$, see, e.g., [18, 10].

If Φ is the equation operator corresponding to either the minimum function, the Fischer-Burmeister function or the penalized Fischer-Burmeister function and x^* is a strongly regular solution of NCP, then it is possible to show that

- (a) all elements $H \in \partial_B \Phi(x^*)$ are nonsingular, and
- (b) Φ satisfies further smoothness assumptions (called *semismoothness/strong semismoothness* in [79]).

These two requirements are exactly those which are needed in order to prove local superlinear or quadratic convergence of the iterates generated by (11), see [79, 78].

Furthermore, it is often relatively easy to make these methods globally convergent at least for monotone problems. To this end, one typically uses the vector

$$d^k := -H_k^{-1} \Phi(x^k)$$

as a search direction and performs a line search along this direction to get sufficient decrease in the corresponding merit function

$$\Psi(x) := \frac{1}{2} \Phi(x)^T \Phi(x). \quad (12)$$

Such a decrease is possible if the merit function is continuously differentiable [18]. This condition is satisfied for the Fischer-Burmeister function and the penalized Fischer-Burmeister function discussed in the previous subsection. In general, these methods work well even for nonmonotone complementarity problems.

Unfortunately, the merit function Ψ corresponding to the minimum function is not everywhere differentiable, and therefore the nonsmooth Newton iteration (11) based on the minimum function is more difficult to globalize. The references [73, 74, 76, 36, 19] give possible strategies.

4.4 Smoothing and Interior-Point Methods

Given an NCP-function ϕ and the corresponding equation operator Φ defined by (10), Proposition 4.8 suggests using nondifferentiable NCP-functions and nonsmooth algorithms. Another possibility to deal with nondifferentiable functions is the smoothing technique whose basis will be described in this section.

Smoothing methods for the NCP typically approximate a nonsmooth NCP-function ϕ by a smooth function ϕ_μ that depends on a smoothing parameter $\mu > 0$. For example, it is possible to approximate the Fischer-Burmeister function by

$$\phi(a, b) = \sqrt{a^2 + b^2} - a - b \approx \sqrt{a^2 + b^2 + \mu} - a - b =: \phi_\mu(a, b). \quad (13)$$

Using similar techniques, it is possible to smooth most known NCP-functions.

As soon as we have such a smooth approximation ϕ_μ , we define the corresponding equation operator

$$\Phi_\mu(x) := \begin{pmatrix} \phi_\mu(x_1, F_1(x)) \\ \vdots \\ \phi_\mu(x_n, F_n(x)) \end{pmatrix}$$

which is now a smooth approximation of Φ if F itself is smooth.

The main idea of any smoothing method is to apply a standard Newton method to the nonlinear system of equations $\Phi_\mu(x) = 0$ and to drive the smoothing parameter μ down to zero. If we denote the value of μ at the k th iteration by μ_k , a typical iteration of a smoothing method is given by

$$x^{k+1} = x^k - \Phi'_{\mu_k}(x^k)^{-1} \Phi_{\mu_k}(x^k), \quad k = 0, 1, 2, \dots$$

A critical point of all smoothing methods is the way the smoothing parameter μ_k gets updated at each iteration. If this is done in an appropriate way, it is possible to prove global convergence for monotone problems, and even local fast convergence if a solution x^* of the NCP satisfies the strong regularity assumption, see [8, 9, 12, 13, 102] for a couple of different smoothing methods.

Smoothing methods have an interesting connection to interior-point methods as pointed out in [44]. For example, let ϕ_μ denote the function defined in (13). Then it is easy to see that

$$\phi_\mu(a, b) = 0 \iff a > 0, b > 0, ab = \mu.$$

Hence solving the nonlinear system of equations

$$\Phi_\mu(x) = 0 \tag{14}$$

is equivalent to solving the central path conditions

$$x_i > 0, \quad F_i(x) > 0, \quad x_i F_i(x) = \mu \quad \forall i = 1, \dots, n \tag{15}$$

which are typically used as the basis of path-following interior-point methods. However, the way smoothing methods and interior-point methods solve their (theoretically equivalent) subproblems (14) and (15), respectively, is completely different. Furthermore, there exist smooth approximations ϕ_μ of NCP-functions ϕ that are not related to the central path conditions, see [11].

So far, the theory developed for interior-point methods (see, e.g., the books [101, 104, 85]) is much stronger than the one developed for smoothing methods. For example, interior-point methods can often be shown to have polynomial complexity and local fast convergence even in the absence of the strong regularity condition. On the other hand, smoothing-type methods seem to be quite promising from a numerical point of view, see [11, 6, 47]. We therefore expect further theoretical improvements in the field of smoothing methods in the near future.

4.5 Linearization Methods

Here, the fundamental idea is to linearize F about the current iterate x^k and solve the corresponding linear complementarity problem

$$x \geq 0, \quad F(x^k) + F'(x^k)(x - x^k) \geq 0, \quad x^T (F(x^k) + F'(x^k)(x - x^k)) = 0 \quad (16)$$

to generate x^{k+1} . In the context of generalized equations, this method, which has been commonly termed SLCP (for sequential LCP), was shown to be locally well-defined and fast convergent under the strong regularity condition [43]. Mathiesen [65, 64] successfully used this scheme for solving equilibrium models such as those outlined in Section 2. While this method works extremely well in the neighborhood of a solution, there are substantial difficulties in globalizing the approach. Two key difficulties are subproblems not having solutions and choosing appropriate merit functions that do not destroy fast local convergence. It is no mere coincidence that these problems plague sequential quadratic programming methods too – their theory can be treated analogously.

For strongly monotone variational inequalities, convergence of line search schemes based on (16) are shown in [34, 97]. A theoretically more general approach is via reformulation of the problem as a nonsmooth system of equations. The normal map [83] defined as

$$F_+(y) := F(y_+) + y - y_+ \quad (17)$$

is equivalent to solving the NCP in the following sense. If y is a zero of (17) then y_+ solves the NCP. Furthermore, if x is a solution of the NCP, then $x - F(x)$ is a zero of the normal map. Note two points that contrast the normal map from previously mentioned nonsmooth equivalents of the NCP. First, it is defined for all points in \mathbb{R}^n even if F is only defined on \mathbb{R}_+^n . It is known to be a piecewise smooth map, the “pieces” correspond to the orthants of \mathbb{R}^n , a subdivided manifold [23], typically called the normal manifold. Secondly, the nonsmoothness is inside the function F as opposed to around the function F , so that F is only evaluated on the positive orthant.

The Newton method proposed in [84], in the context of the normal map, approximates $F_+(y)$ with a piecewise linear normal map

$$A^k(y) := F(y_+^k) + F'(y_+^k)(y_+ - y_+^k) + y - y_+.$$

The Newton point is a zero of A^k . It is well known that this is equivalent to SLCP by taking $x^k = y_+^k$. However, in [80], a globalization scheme was proposed that extends damped Newton methods for smooth equations to the above situation. Instead of using Lemke’s method [16, 51] to solve (16), a path $p^k(t)$ is constructed [21] from y^k to y^{k+1} with the property that

$$A^k(p^k(t)) = (1 - t)F_+(y^k).$$

This path has exactly the same properties that are required for standard line search methods, namely that small steps guarantee sufficient decrease. That theory can be extended to this case, provided that $\|F_+(y)\|$ is used as the merit function, to prove standard global and local convergence results [21, 80].

Considerable success has been reported for an implementation of this approach called the PATH solver [21, 30]. This implementation uses standard large scale linear programming (simplex) technology [69] to generate the required path. SLCP methods typically use similar technology, except the subproblems use ray starts in Lemke's method as opposed to the required regular point starts in PATH. Some advanced techniques for choosing the initial piece of the normal manifold can be found in [1, 22].

Most of the above implementations rely on heuristics to deal with the case where the subproblems have no solution. Recent work in this area has extended the global convergence theory along different lines. A theoretical framework for globalization of feasible descent methods was proposed in [28] that uses (8) as a merit function. The main features of this framework are: (a) it is well-defined for an arbitrary complementarity problem, (b) it generates only feasible iterates, (c) it has a strong global convergence theory, and (d) it is locally fast convergent under standard regularity assumptions. The key observations of such a globalization are that if the subproblems are unsolvable, then a projected gradient step can be taken for the merit function, guaranteeing progress and that eventually the full Newton step will be accepted. Numerical results in the cited paper indicate this framework leads to substantial computational improvements.

5 Extensions

5.1 Mixed Complementarity Problems

While most of this paper has been written in the context of nonlinear complementarity problems, many practical applications generate problems for which some of the variables are nonnegative, others are bounded, and others are completely free. To accommodate these problems, many researchers have investigated the mixed complementarity problem (MCP), of finding a vector $x \in [l, u]$ such that exactly one of the following holds:

$$\begin{aligned} x_i = l_i & \quad \text{and} \quad F_i(x) > 0, \\ x_i = u_i & \quad \text{and} \quad F_i(x) < 0, \\ x_i \in [l_i, u_i] & \quad \text{and} \quad F_i(x) = 0. \end{aligned}$$

Note that l_i and u_i can be infinite. Two special cases are of importance. When $l_i = -\infty$ and $u_i = \infty$ then the MCP is exactly the problem of finding a zero of F . This shows why many of the techniques for solving complementarity problems are closely related to nonlinear equation theory, as opposed to standard optimization techniques. Furthermore, when $l_i = 0$ and $u_i = \infty$ the NCP is recovered, and thus it is clear that MCP is a generalization of NCP. Although there are many computational issues related to exploiting this framework, it is well known that appropriate changes of variables can reduce the MCP to a (larger dimensioned) NCP. Therefore, for theoretical purposes it is entirely adequate to deal exclusively with NCP's. In practice, however, exploiting the MCP structure is of paramount importance [20, 32].

5.2 Variational Inequalities

Given a nonempty, closed and convex subset $X \subseteq \mathbb{R}^n$ and a function $F : X \rightarrow \mathbb{R}^n$, the variational inequality problem (VIP) is to find a vector $x^* \in \mathbb{R}^n$ such that

$$F(x^*)^T(x - x^*) \geq 0 \quad \text{for all } x \in X.$$

It is not difficult to see that the variational inequality problem VIP reduces to NCP if the feasible set X is equal to the nonnegative orthant \mathbb{R}_+^n , and becomes the MCP if $X = [l, u]$.

Many results and algorithms described above for the complementarity problem can actually be extended to the more general variational inequality problem. For example, Theorem 3.3 still holds if NCP is replaced everywhere by VIP. Also Theorem 3.5 has an analogous result for variational inequalities [39].

For solution techniques, all the projection-type methods outlined in Section 4.1 can be applied to VIP and have the same global convergence properties if we interpret the symbol z_+ as the Euclidean projection of a vector $z \in \mathbb{R}^n$ onto the feasible set X (rather than the projection onto \mathbb{R}_+^n).

Also the linearization methods have a direct translation for variational inequalities. For example, the SLCP method for VIP generates a sequence $\{x^k\}$ in such a way that x^{k+1} is a solution of the linearized variational inequality

$$(F(x^k) + F'(x^k)(x - x^k))^T (y - x) \geq 0 \quad \text{for all } y \in X,$$

with similar convergence to that described in Section 4.5.

Furthermore, there exist a couple of gap function approaches for the solution of VIP which may be viewed as a generalization of the merit function approaches used for complementarity problems although the situation is much more complicated for VIP.

The *gap function* was introduced in [4] and is defined by

$$g(x) := \sup_{y \in X} \{F(x)^T(x - y)\}.$$

It is not difficult to see that g is nonnegative on the feasible set X and that $g(x) = 0$ and $x \in X$ if and only if x solves VIP. Hence VIP can be rewritten as the constrained minimization problem

$$\min g(x) \quad \text{s.t.} \quad x \in X.$$

Some algorithms based on this are given in [61, 63]. However, the gap function g is not differentiable in general. Moreover, when X is unbounded, g is in general not finite-valued.

These drawbacks can be avoided by using the *regularized gap function* [3, 34]

$$f_\alpha(x) := \max_{y \in X} \left\{ F(x)^T(x - y) - \frac{\alpha}{2} \|x - y\|^2 \right\},$$

where α is any positive parameter. It is also nonnegative on the set X and x solves VIP if and only if $f_\alpha(x) = 0$ and $x \in X$. Hence, VIP is equivalent to the constrained optimization problem

$$\min f_\alpha(x) \quad \text{s.t.} \quad x \in X. \tag{18}$$

In addition, f_α can be shown to be continuously differentiable and finite-valued everywhere.

However, the regularized gap function has a serious drawback: If x^* denotes a constrained stationary point of (18), then F has to be strongly monotone in order to guarantee that x^* is a solution of VIP. Another extension, the *D-gap function* [77, 103] also suffers from the same drawback.

5.3 Mathematical Programs with Equilibrium Constraints

A mathematical program with equilibrium constraints (MPEC) is a constrained optimization problem with some of the constraints representing certain equilibrium conditions. For example, if these equilibrium conditions are described by a complementarity problem, then an MPEC can be formulated as

$$\min f(x, y) \quad \text{s.t.} \quad x \in X, y \geq 0, F(x, y) \geq 0, y^T F(x, y) = 0; \quad (19)$$

here, $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ and $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ are given functions and $X \subseteq \mathbb{R}^n$ is an abstract constraint which can usually be represented by standard equality/inequality constraints.

Note that the MPEC (19) has two types of variables, the design variables x and the state variables y . The complementarity constraints in (19) are parameterized in the design variables. While extensions or particular instantiations of (19) are possible, we will restrict ourselves to this form in order to convey the main algorithmic ideas.

The first point is that (19) should not be treated as a standard constrained optimization problem, since it is well known that it does not even satisfy the Mangasarian-Fromovitz constraint qualification [14]. Therefore, standard methods from constrained optimization are prone to failure.

Several researchers have proposed more specialized solution methods for MPEC problems. Given there is much current research in this area, we will only summarize a few central ideas.

One way to overcome the previously mentioned difficulty in (19) is to add the constraint $y^T F(x, y) = 0$ to the objective function and then to solve a sequence of penalized problems

$$\min f(x, y) + \rho y^T F(x, y) \quad \text{s.t.} \quad x \in X, y \geq 0, F(x, y) \geq 0$$

with penalty parameter $\rho > 0$. In general, this gives only an inexact penalty reformulation unless some strong conditions hold [52, 58, 89, 90]. Some researchers also proposed exact penalty functions for the MPEC (19) which, however, are severely nondifferentiable and therefore difficult to deal with from a numerical point of view.

Another technique for the solution of the MPEC (19) is the implicit programming approach. These methods assume that, for any given $x \in X$, one can find a solution $y(x)$ of the complementarity constraints in (19). They replace the variable y by the function $y(x)$ in the objective function f and try to solve the optimization problem

$$\min f(x, y(x)) \quad \text{s.t.} \quad x \in X \quad (20)$$

which is a problem in the design variables only. The function $y(x)$ is usually nondifferentiable, so one has to use nondifferentiable optimization methods to solve (20), see [71, 72].

Another possibility to deal with the MPEC (19) is to reformulate the complementarity constraints using an NCP-function ϕ . One obtains a constrained optimization problem of the form

$$\min f(x, y) \quad \text{s.t.} \quad x \in X, \phi(y_i, F_i(x, y)) = 0 \quad (i = 1, \dots, m).$$

In order to avoid the nonsmooth constraints $\phi(y_i, F_i(x, y)) = 0$, we can use the smoothing idea and approximate this problem by the smooth optimization problem

$$\min f(x, y) \quad \text{s.t.} \quad x \in X, \phi_\mu(y_i, F_i(x, y)) = 0 \quad (i = 1, \dots, m),$$

where, of course, ϕ_μ is a smooth approximation for ϕ , see e.g., (13). This idea has been exploited in [25, 35, 42]. Alternatively, and closely related to the smoothing idea, we can apply interior-point techniques to the perturbed problem

$$\min f(x, y) \quad \text{s.t.} \quad x \in X, y_i > 0, F_i(x, y) > 0, y_i F_i(x, y) = \mu \quad (i = 1, \dots, m).$$

For some further techniques and several theoretical results about MPECs, we refer the reader to [52, 70].

References

- [1] K. M. Anstreicher, J. Lee, and T. F. Rutherford. Crashing a maximum-weight complementarity basis. *Mathematical Programming*, 54:281–294, 1992.
- [2] K. Arrow and G. Debreu. Existence of equilibrium for a competitive economy. *Econometrica*, 22:265–290, 1954.
- [3] G. Auchmuty. Variational principles for variational inequalities. *Numerical Functional Analysis and Optimization*, 10:863–874, 1989.
- [4] A. Auslender. *Optimization Méthodes Numériques*. Masson, Paris, 1976.
- [5] D. P. Bertsekas and J. N. Tsitsiklis. *Parallel and Distributed Computation: Numerical Methods*. Prentice-Hall, Inc, Englewood Cliffs, New Jersey, 1989.
- [6] S. C. Billups, S. P. Dirkse, and M. C. Ferris. A comparison of large scale mixed complementarity problem solvers. *Computational Optimization and Applications*, 7:3–25, 1997.
- [7] G. Bolzon, G. Maier, and F. Tin-Loi. Holonomic and nonholonomic simulations of quasi-brittle fracture: a comparative study of mathematical programming approaches. In F. H. Wittmann, editor, *Fracture Mechanics of Concrete Structures*, Second International Conference on Fracture Mechanics of Concrete Structures (FRAMCOS 2), pages 885–898. Aedificatio Publishers, 1995.
- [8] J. V. Burke and S. Xu. The global linear convergence of a non-interior path-following algorithm for linear complementarity problems. *Mathematics of Operations Research*, 23:719–735, 1998.

- [9] B. Chen and X. Chen. A global and local superlinear continuation-smoothing method for $p_0 + r_0$ and monotone NCP. *SIAM Journal on Optimization*, forthcoming, 1998.
- [10] B. Chen, X. Chen, and C. Kanzow. A penalized Fischer-Burmeister NCP-function: Theoretical investigation and numerical results. Preprint 126, Institute of Applied Mathematics, University of Hamburg, Hamburg, Germany, 1997.
- [11] Chunhui Chen and O. L. Mangasarian. A class of smoothing functions for nonlinear and mixed complementarity problems. *Computational Optimization and Applications*, 5:97–138, 1996.
- [12] X. Chen, L. Qi, and D. Sun. Global and superlinear convergence of the smoothing Newton method and its application to general box constrained variational inequalities. *Mathematics of Computation*, forthcoming, 1998.
- [13] X. Chen and Y. Ye. On homotopy-smoothing methods for variational inequalities. *SIAM Journal on Control and Optimization*, forthcoming, 1998.
- [14] Y. Chen and M. Florian. The nonlinear bilevel programming problem: Formulations, regularity, and optimality conditions. *Optimization*, 32:193–209, 1995.
- [15] F. H. Clarke. *Optimization and Nonsmooth Analysis*. John Wiley & Sons, New York, 1983.
- [16] R. W. Cottle and G. B. Dantzig. Complementary pivot theory of mathematical programming. *Linear Algebra and Its Applications*, 1:103–125, 1968.
- [17] R. W. Cottle, J. S. Pang, and R. E. Stone. *The Linear Complementarity Problem*. Academic Press, Boston, 1992.
- [18] T. De Luca, F. Facchinei, and C. Kanzow. A semismooth equation approach to the solution of nonlinear complementarity problems. *Mathematical Programming*, 75:407–439, 1996.
- [19] T. De Luca, F. Facchinei, and C. Kanzow. A theoretical and numerical comparison of some semismooth algorithms for complementarity problems. Mathematical Programming Technical Report 97-15, Computer Sciences Department, University of Wisconsin, Madison, Wisconsin, 1997.
- [20] S. P. Dirkse and M. C. Ferris. MCPLIB: A collection of nonlinear mixed complementarity problems. *Optimization Methods and Software*, 5:319–345, 1995.
- [21] S. P. Dirkse and M. C. Ferris. The PATH solver: A non-monotone stabilization scheme for mixed complementarity problems. *Optimization Methods and Software*, 5:123–156, 1995.
- [22] S. P. Dirkse and M. C. Ferris. Crash techniques for large-scale complementarity problems. In Ferris and Pang [31], pages 40–61.

- [23] B. C. Eaves. A short course in solving equations with PL homotopies. In R. W. Cottle and C. E. Lemke, editors, *Nonlinear Programming*, pages 73–143, Providence, RI, 1976. American Mathematical Society, SIAM–AMS Proceedings.
- [24] Y. G. Evtushenko and V. A. Purtov. Sufficient conditions for a minimum for nonlinear programming problems. *Soviet Mathematics Doklady*, 30:313–316, 1984.
- [25] F. Facchinei, H. Jiang, and L. Qi. A smoothing method for mathematical programs with equilibrium constraints. *Mathematical Programming*, forthcoming, 1998.
- [26] F. Facchinei and J. Soares. A new merit function for nonlinear complementarity problems and a related algorithm. *SIAM Journal on Optimization*, 7:225–247, 1997.
- [27] M. C. Ferris, R. Fourer, and D. M. Gay. Expressing complementarity problems and communicating them to solvers. Mathematical Programming Technical Report 98-02, Computer Sciences Department, University of Wisconsin, Madison, Wisconsin, 1998.
- [28] M. C. Ferris, C. Kanzow, and T. S. Munson. Feasible descent algorithms for mixed complementarity problems. Mathematical Programming Technical Report 98-04, Computer Sciences Department, University of Wisconsin, Madison, Wisconsin, 1998.
- [29] M. C. Ferris and T. S. Munson. Complementarity problems in GAMS and the PATH solver. Mathematical Programming Technical Report 98-12, Computer Sciences Department, University of Wisconsin, Madison, Wisconsin, 1998.
- [30] M. C. Ferris and T. S. Munson. Interfaces to PATH 3.0: Design, implementation and usage. *Computational Optimization and Applications*, forthcoming, 1998.
- [31] M. C. Ferris and J. S. Pang, editors. *Complementarity and Variational Problems: State of the Art*, Philadelphia, Pennsylvania, 1997. SIAM Publications.
- [32] M. C. Ferris and J. S. Pang. Engineering and economic applications of complementarity problems. *SIAM Review*, 39:669–713, 1997.
- [33] A. Fischer. A special Newton–type optimization method. *Optimization*, 24:269–284, 1992.
- [34] M. Fukushima. Equivalent differentiable optimization problems and descent methods for asymmetric variational inequality problems. *Mathematical Programming*, 53:99–110, 1992.
- [35] M. Fukushima, Z.-Q. Luo, and J. S. Pang. A globally convergent sequential quadratic programming algorithm for mathematical programs with linear complementarity constraints. *Computational Optimization and Applications*, 10:5–34, 1998.
- [36] S. A. Gabriel and J. S. Pang. An inexact NE/SQP method for solving the nonlinear complementarity problem. *Computational Optimization and Applications*, 1:67–91, 1992.

- [37] C. Geiger and C. Kanzow. On the resolution of monotone complementarity problems. *Computational Optimization and Applications*, 5:155–173, 1996.
- [38] P. T. Harker. *Lectures on Computation of Equilibria with Equation-Based Methods*. CORE Lecture Series. CORE Foundation, Louvain-la-Neuve, Université Catholique de Louvain, 1993.
- [39] P. T. Harker and J. S. Pang. Finite-dimensional variational inequality and nonlinear complementarity problems: A survey of theory, algorithms and applications. *Mathematical Programming*, 48:161–220, 1990.
- [40] G. W. Harrison, T. F. Rutherford, and D. Tarr. Quantifying the Uruguay round. *Economic Journal*, 107, 1997.
- [41] A. N. Iusem and B. F. Svaiter. A variant of Korpelevich’s method for variational inequalities with a new search strategy. *Optimization*, 42:309–321, 1997.
- [42] H. Jiang and D. Ralph. Smooth SQP methods for mathematical programs with nonlinear complementarity constraints. Technical report, Department of Mathematics and Statistics, The University of Melbourne, Australia, 1997.
- [43] N. H. Josephy. Newton’s method for generalized equations. Technical Summary Report 1965, Mathematics Research Center, University of Wisconsin, Madison, Wisconsin, 1979.
- [44] C. Kanzow. Some noninterior continuation methods for linear complementarity problems. *SIAM Journal on Matrix Analysis and Applications*, 17:851–868, 1996.
- [45] C. Kanzow and H. Kleinmichel. A class of Newton-type methods for equality and inequality constrained optimization. *Optimization Methods and Software*, 5:173–198, 1995.
- [46] C. Kanzow and H. Kleinmichel. A new class of semismooth Newton-type methods for nonlinear complementarity problems. *Computational Optimization and Applications*, forthcoming, 1998.
- [47] C. Kanzow and H. Pieper. Jacobian smoothing methods for nonlinear complementarity problems. *SIAM Journal on Optimization*, forthcoming, 1998.
- [48] C. Kanzow, N. Yamashita, and M. Fukushima. New NCP-functions and their properties. *Journal of Optimization Theory and Applications*, 94:115–135, 1997.
- [49] E. N. Khobotov. A modification of the extragradient method for the solution of variational inequalities and some optimization problems. *Zhurnal Vychislitel’noi Matematiki i Matematicheskoi Fiziki*, 27:1462–1473, 1987.
- [50] G. M. Korpelevich. The extragradient method for finding saddle points and other problems. *Matecon*, 12:747–756, 1976.

- [51] C. E. Lemke and J. T. Howson. Equilibrium points of bimatrix games. *SIAM Journal on Applied Mathematics*, 12:413–423, 1964.
- [52] Z.-Q. Luo, J. S. Pang, and D. Ralph. *Mathematical Programs with Equilibrium Constraints*. Cambridge University Press, 1996.
- [53] Z.-Q. Luo and P. Tseng. A new class of merit functions for the nonlinear complementarity problem. In Ferris and Pang [31], pages 204–225.
- [54] G. Maier. A matrix structural theory of piecewise linear elastoplasticity with interacting yield planes. *Meccanica*, 5:54–66, 1970.
- [55] G. Maier. Incremental plastic analysis in the presence of large displacement and physical instabilizing effects. *International Journal of Solids and Structures*, 7:345–372, 1971.
- [56] G. Maier and G. Novati. A shakedown and bounding theory allowing for nonlinear hardening and second order geometric effects with reference to discrete structural models. In *Inelastic Solids and Structures, A. Sawczuk memorial volume*, pages 451–471. Pineridge Press, Swansea, 1990.
- [57] O. L. Mangasarian. Equivalence of the complementarity problem to a system of nonlinear equations. *SIAM Journal on Applied Mathematics*, 31:89–92, 1976.
- [58] O. L. Mangasarian and J. S. Pang. Exact penalty functions for mathematical programs with linear complementarity constraints. *Optimization*, 42:1–8, 1997.
- [59] O. L. Mangasarian and M. V. Solodov. Nonlinear complementarity as unconstrained and constrained minimization. *Mathematical Programming*, 62:277–298, 1993.
- [60] A. S. Manne and T. F. Rutherford. International trade in oil, gas and carbon emission rights: An intertemporal general equilibrium model. *The Energy Journal*, 14:1–20, 1993.
- [61] P. Marcotte. A new algorithm for solving variational inequalities with application to the traffic assignment problem. *Mathematical Programming*, 33:339–351, 1985.
- [62] P. Marcotte. Application of Khobotov’s algorithm to variational inequalities and network equilibrium problems. *Information Systems and Operational Research*, 29:258–270, 1991.
- [63] P. Marcotte and J.-P. Dussault. A note on a globally convergent Newton method for solving monotone variational inequalities. *Operations Research Letters*, 6:35–42, 1987.
- [64] L. Mathiesen. Computation of economic equilibria by a sequence of linear complementarity problems. *Mathematical Programming Study*, 23:144–162, 1985.

- [65] L. Mathiesen. An algorithm based on a sequence of linear complementarity problems applied to a Walrasian equilibrium model: An example. *Mathematical Programming*, 37:1–18, 1987.
- [66] J. J. Moré and W. C. Rheinboldt. On P- and S-functions and related classes of n-dimensional nonlinear mappings. *Linear Algebra and Its Applications*, 6:45–68, 1973.
- [67] A. Nagurney. *Network Economics – A Variational Inequality Approach*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993.
- [68] J. F. Nash. Equilibrium points in N-person games. *Proceedings of the National Academy of Sciences*, 36:48–49, 1950.
- [69] J. L. Nazareth. *Computer Solution of Linear Programs*. Oxford University Press, Oxford, 1987.
- [70] J. Outrata, M. Kočvara, and J. Zowe. *Nonsmooth Approach to Optimization Problems with Equilibrium Constraints*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1998.
- [71] J. V. Outrata. On optimization problems with variational inequality constraints. *SIAM Journal on Optimization*, 4:340–357, 1994.
- [72] J. V. Outrata and J. Zowe. A numerical approach to optimization problems with variational inequality constraints. *Mathematical Programming*, 68:105–130, 1995.
- [73] J. S. Pang. Newton’s method for B-differentiable equations. *Mathematics of Operations Research*, 15:311–341, 1990.
- [74] J. S. Pang. A B-differentiable equation based, globally and locally quadratically convergent algorithm for nonlinear programs, complementarity and variational inequality problems. *Mathematical Programming*, 51:101–132, 1991.
- [75] J. S. Pang. Complementarity problems. In R. Horst and P. Pardalos, editors, *Handbook in Global Optimization*. Kluwer Academic Publishers, Boston, 1994.
- [76] J. S. Pang and S. A. Gabriel. NE/SQP: A robust algorithm for the nonlinear complementarity problem. *Mathematical Programming*, 60:295–338, 1993.
- [77] J. M. Peng. Equivalence of variational inequality problems to unconstrained optimization. *Mathematical Programming*, 78:347–356, 1997.
- [78] L. Qi. Convergence analysis of some algorithms for solving nonsmooth equations. *Mathematics of Operations Research*, 18:227–244, 1993.
- [79] L. Qi and J. Sun. A nonsmooth version of Newton’s method. *Mathematical Programming*, 58:353–368, 1993.
- [80] D. Ralph. Global convergence of damped Newton’s method for nonsmooth equations, via the path search. *Mathematics of Operations Research*, 19:352–389, 1994.

- [81] S. M. Robinson. Strongly regular generalized equations. *Mathematics of Operations Research*, 5:43–62, 1980.
- [82] S. M. Robinson. Local structure of feasible sets in nonlinear programming, Part III: Stability and sensitivity. *Mathematical Programming Study*, 30:45–66, 1987.
- [83] S. M. Robinson. Normal maps induced by linear transformations. *Mathematics of Operations Research*, 17:691–714, 1992.
- [84] S. M. Robinson. Newton’s method for a class of nonsmooth functions. *Set Valued Analysis*, 2:291–305, 1994.
- [85] C. Roos, T. Terlaky, and J.-Ph. Vial. *Theory and Algorithms for Linear Optimization: An Interior-Point Approach*. John Wiley & Sons, Chichester, 1997.
- [86] T. F. Rutherford. Extensions of GAMS for complementarity problems arising in applied economic analysis. *Journal of Economic Dynamics and Control*, 19:1299–1324, 1995.
- [87] T. F. Rutherford. Applied general equilibrium modeling with MPSGE as a GAMS subsystem: An overview of the modeling framework and syntax. *Computational Economics*, forthcoming, 1998.
- [88] H. E. Scarf. *The Computation of Economic Equilibria*. Yale University Press, New Haven, Connecticut, 1973.
- [89] H. Scheel and S. Scholtes. Mathematical programs with complementarity constraints: Stationarity, optimality, and sensitivity. Technical report, Judge Institute of Management Studies, University of Cambridge, Cambridge, England, 1998.
- [90] S. Scholtes and M. Stöhr. Exact penalization of mathematical programs with equilibrium constraints. *SIAM Journal on Control and Optimization*, forthcoming, 1998.
- [91] M. Sibony. Methodes iteratives pur les equations et inequations aux derives partielles nonlineaires de type monotone. *Calcolo*, 7:65–183, 1970.
- [92] Y. Smeers. Computable equilibrium models and the restructuring of the European electricity and gas markets. *The Energy Journal*, 1997.
- [93] M. V. Solodov and B. F. Svaiter. A new projection method for variational inequality problems. *SIAM Journal on Control and Optimization*, forthcoming, 1998.
- [94] M. V. Solodov and P. Tseng. Modified projection–type methods for monotone variational inequalities. *SIAM Journal on Control and Optimization*, 34:1814–1830, 1996.
- [95] D. Sun. A class of iterative methods for nonlinear projection equations. *Journal of Optimization Theory and Applications*, 91:123–140, 1996.
- [96] D. Sun and L. Qi. On NCP-functions. *Computational Optimization and Applications*, forthcoming, 1998.

- [97] K. Taji, M. Fukushima, and T. Ibaraki. A globally convergent Newton method for solving strongly monotone variational inequalities. *Mathematical Programming*, 58:369–383, 1993.
- [98] F. Tin-Loi and M. C. Ferris. Holonomic analysis of quasibrittle fracture with nonlinear softening. In B. L. Karihaloo, Y. W. Mai, M. I. Ripley, and R. O. Ritchie, editors, *Advances in Fracture Research*, volume 2, pages 2183–2191. Pergamon Press, 1997.
- [99] F. Tin-Loi and J. S. Misa. Large displacement elastoplastic analysis of semirigid steel frames. *International Journal for Numerical Methods in Engineering*, 39:741–762, 1996.
- [100] F. Tin-Loi and J. S. Pang. Elastoplastic analysis of structures with nonlinear hardening. *Computer Methods in Applied Mechanics and Engineering*, 107:299–312, 1993.
- [101] S. J. Wright. *Primal–Dual Interior–Point Methods*. SIAM, Philadelphia, Pennsylvania, 1997.
- [102] S. Xu and J. V. Burke. A polynomial time interior-point path-following algorithm for LCP based on Chen-Harker-Kanzow smoothing techniques. *Mathematical Programming*, forthcoming, 1998.
- [103] N. Yamashita, K. Taji, and M. Fukushima. Unconstrained optimization reformulations of variational inequality problems. *Journal of Optimization Theory and Applications*, 92:439–456, 1997.
- [104] Y. Ye. *Interior-Point Algorithms: Theory and Analysis*. John Wiley & Sons, New York, NY, 1997.