

**A CHARACTERIZATION OF THE APPROXIMATION
ORDER OF MULTIVARIATE SPLINE SPACES**

by

Amos Ron

Computer Sciences Technical Report #885

November 1989

UNIVERSITY OF WISCONSIN-MADISON
CENTER FOR THE MATHEMATICAL SCIENCES

A characterization of the approximation order of multivariate spline spaces

Amos Ron[†]

Center for the Mathematical Sciences
and
Computer Sciences Department
1210 W. Dayton St.
University of Wisconsin-Madison
Madison, WI 53706

November 1989

ABSTRACT

We analyze the approximation order associated with a directed set of spaces, $\{S_h\}_{h>0}$, each of which spanned by the $h\mathbb{Z}^s$ -translates of one compactly supported function $\phi_h : \mathbb{R}^s \rightarrow \mathbb{C}$. Under a regularity condition on the sequence $\{\phi_h\}_h$, we show that the optimal approximation order (in the ∞ -norm) is always realized by quasi-interpolants, hence in a linear way. These quasi-interpolants provide the best approximation rates from $\{S_h\}_h$ to an exponential space of good approximation order at the origin.

As for the case when each S_h is obtained by scaling S_1 , under the assumption

$$\sum_{\alpha \in \mathbb{Z}^s} \phi_1(\cdot - \alpha) \neq 0, \tag{*}$$

the results here provide an unconditional characterization of the best approximation order in terms of the polynomials in S_1 . The necessity of (*) in this characterization is demonstrated by a counterexample.

AMS (MOS) Subject Classifications: primary 41A15, 41A25, 41A40, 41A63; secondary 65D15.

Key Words and phrases: approximation order, Strang-Fix Conditions, exponentials, polynomials, multivariate, multivariate splines, uniform mesh, regular grids, integer translates, quasi-interpolation.

[†] Supported in part by Carl de Boor's Steenbock Professorship, University of Wisconsin-Madison.

A characterization of the approximation order of multivariate spline spaces

Amos Ron

1. Introduction

The determination of the approximation order associated with spline spaces is one of the major and fundamental goals of spline theory. In an abstract form, we are given a collection of function spaces $\{S_h\}_h$ where h is a continuous or discrete positive parameter, and look for the largest d for which

$$\text{dist}(f, S_h) = O(h^d), \quad \forall f \in F,$$

where dist is measured by some metric or norm (usually a p - or Sobolev norm) and F is an admissible function space with respect to dist . Here, we always assume that all functions are complex-valued and defined on the vector space \mathbb{R}^s . We think of h as measuring (in a linear way) the roughness of the mesh used, hence indeed expect to approximate better as $h \rightarrow 0$. A particular case of interest, which will be referred to as “**the scaling case**” occurs when the refined spaces S_h are dilations of S_1 , i.e.,

$$S_h = \sigma_{1/h} S_1,$$

with σ_h the scaling operator

$$\sigma_h : f \mapsto f(h \cdot).$$

In multivariate splines, as well as in finite elements, one prefers approximant spaces which are spanned by compactly supported functions, since then there is hope to provide a **local** scheme for approximation from the underlying space. In the area of **multivariate splines on regular meshes**, one also assumes that the spaces S_h are translation-invariant, or more precisely, each S_h is invariant under $h\mathbb{Z}^s$ -translates. This leads to the model where each of the spaces S_h is assumed to be spanned by the $h\mathbb{Z}^s$ -translates of one or finitely many compactly supported functions. In general, it is desirable to avoid the assumption that these translates are linearly independent, i.e., form a basis for S_h , since such condition is not met in many cases of practical interest (e.g., the Zwart element and also most of the cases when more than one compactly supported function is involved).

The simplest case therefore occurs when we assume both, that S_1 is spanned by the integer translates of a single compactly supported function ϕ and that S_h is the h -scale of S_1 hence is spanned by the h -scale of ϕ . This model was investigated intensively in the late 60's-early 70's by people in the finite element area. The following theorem is essentially Theorem 1 of [SF] (at least for $p = 2, \infty$) and these days is usually referred to as **The Strang-Fix Conditions**. In the statement of the theorem we use the notation $\phi *_h$ for the semi-discrete convolution operator

$$\phi *_h : \mathbb{C}^{h\mathbb{Z}^s} \rightarrow S_h, \quad \phi * : a \mapsto \sum_{\alpha \in h\mathbb{Z}^s} a(\alpha) \phi(\cdot - \alpha), \quad \phi * := \phi *_1,$$

and $\phi *_h'$ for the semi-discrete operator

$$\phi *_h' f := \phi *_h (f|_{h\mathbb{Z}^s}), \quad \phi *_1' := \phi *_1'.$$

Also, π_k stands for the space of polynomials of (total) degree at most k .

(1.1) **Theorem.** Let ϕ be a compactly supported continuous function. Let S_1 be the space spanned by the integer translates of ϕ and $S_h := \sigma_{1/h} S_1$. Then, for $1 \leq p \leq \infty$ the following conditions are equivalent

- (1) $\hat{\phi}(0) \neq 0$, and for $|\alpha| \leq d-1$, $D^\alpha \hat{\phi}$ vanishes on $2\pi\mathbb{Z}^s \setminus 0$.
- (2) For every $p \in \pi_{d-1}$, $\phi *' p$ is a polynomial of the form $cp + q$, for some $c \neq 0$ and $q \in \pi_{<\deg p}$.
- (3) For every smooth function f which is in L_p together with all its derivatives up to order d , there exist sequences $a_h : h\mathbb{Z}^s$ satisfying
 - (3a) $\|f - \sigma_{1/h} \phi *'_h a_h\|_p = O(h^d)$.
 - (3b) $\|a_h\|_{\ell_p} \leq kh^{-s/p} \|f\|_p$.

The equivalence of (1) and (2) was refined and extended since then by several authors (cf. e.g., [DM₁] and [B₁]; furthermore, [B₁] shows that the two first conditions are equivalent to the apparently weaker one: $\hat{\phi}(0) \neq 0$, $\pi_{d-1} \subset S_1$), yet is mainly technical since it deals with a characterization of the **polynomials** in S_1 in terms of the Fourier transform $\hat{\phi}$ and is not related directly to the **approximation order**. The implication (1,2) \implies (3) initiated much work by many authors aiming at constructing explicit linear maps into S_h , the so-called quasi-interpolants, which satisfy the conditions in (3), and which (by now) are quite well understood (cf. [B₂] for an updated survey of quasi-interpolants). Still, it was probably the saturation result of the above theorem that was so spectacular: one cannot achieve approximation order d from the dilated spaces $\{S_h\}_h$ unless $\pi_{d-1} \subset S_1$! However, one should bear in mind that condition (3) is assuming a restrictive way for approximation from S_h (termed “controlled” by Dahmen and Micchelli), hence does not characterize the approximation order from $\{S_h\}_h$ for smooth functions, and gives only a **lower bound** for that approximation order. As a matter of fact the desired approximation order from spaces of the above form is known only in some special cases (such as when ϕ is a box spline, [BH₁]). In their way to generalize the Strang-Fix Conditions from the case of a singleton to the case of several ϕ ’s, de Boor and Jia [BJ], introduced a different notion of constrained approximation (which they referred to as “local approximation”) which allows the extension of the above saturation result to the case of several functions; this still left open the question of the approximation order itself. Replacing the “controlled” notion by another constrained approximation notion introduced in [BJ] (and referred there to as “local approximation”), (1.1)Theorem admits a generalization to the case of several compactly supported functions, [SF], [BJ], however still leaving open the question of the unconditional approximation order. Finally, the analysis in [DR] (of the approximation order for exponential box spline spaces) demonstrated the fact that good rates of approximation can be obtained from spaces $\{S_h\}$, where each one of them is spanned as before by the translates of a single function, yet none of them contains non-trivial polynomials. (Clearly, such spaces are **not** obtained by dilating S_1). In view of these latter results, one may interpret the Strang-Fix Conditions as saying that constructing S_h by scaling S_1 is appropriate only when S_1 contains a sufficiently large space of polynomials.

In this paper, we characterize the **order of best approximation** from spaces of the form $\{S_h := \sigma_{1/h} S_1\}_h$, with S_1 spanned by the integer translates of a compactly supported function ϕ .

We show that, under the assumption

$$\sum_{\alpha \in \mathbb{Z}^s} \phi(\cdot - \alpha) \neq 0,$$

the approximation order associated with $\{S_h\}_h$ is determined by the polynomials in S_1 exactly in the same way the controlled and local approximation orders are characterized, thus improving the statement of Strang-Fix Conditions above. Our course though, does not focus on (and as a matter of fact is not aimed at) spaces obtained by scalings, but considers the general setting of an arbitrary directed function space set $\{S_h\}_h$, each of which is spanned by the $h\mathbb{Z}^s$ -translates of a compactly supported function ϕ_h . The goal is to provide information about the approximation order of $\{S_h\}_h$ and about maps that realize this approximation order. In order to make a good use of the compact support assumption, it is essential to assume that for some $k > 0$

$$\text{supp } \phi_h \subset [0, hk]^s, \forall h,$$

since otherwise the translates of ϕ_h are not uniformly local as $h \rightarrow 0$.

A major step in the characterization here of the above mentioned approximation order is the construction, for a given finite dimensional space H of a certain type, a sequence $\{Q_h\}_h$ of uniformly local, uniformly bounded quasi-interpolants (each maps into the associated S_h), that approximates the elements of H to the same order as best approximations do. In that process H is required to be a translation-invariant subspace of L_∞ , which leads naturally to exponential spaces H spanned by exponentials of imaginary frequencies. Using quasi-interpolation arguments, we show that the operators $\{Q_h\}_h$ provide approximants of optimal order to **all** admissible functions, in case H approximates well locally (say, at the origin). This gives rise to a scheme for the determination of the approximation order from $\{S_h\}_h$, in which finding this order is reduced to determining the approximation order obtained by applying a prescribed set of quasi-interpolants to prescribed (and finitely many) exponentials. This also proves that the best approximation order can always be realized in a linear way (i.e., by a quasi-interpolant), and furthermore, that we can construct quasi-interpolants of optimal order (essentially) without an a priori (or even a posteriori) knowledge about this approximation order.

In the special case when $\phi_h = \sigma_{1/h}\phi_1$, the quasi-interpolants constructed are not scales one of the other (i.e., $Q_h \neq \sigma_{1/h}Q_1\sigma_h$), yet it is shown that the scaled operators $\{\sigma_h Q_h\}_h$ **converge**, and the limiting quasi-interpolant Q is then used to obtain the mentioned above improvement of the Strang-Fix Conditions.

In section 2 we analyze first the relatively simpler case of approximation order to smooth bounded functions, an analysis which serves as an illustration of the general case, while being better tight in its statements and error estimates. The main results together with their proofs are given in section 3. Finally, in section 4 we consider the scaling case under the singularity assumption $\sum_{\alpha \in \mathbb{Z}^s} \phi(\alpha) = 0$. We prove a theorem which allows us to embed (in a suitable sense) the case of several ϕ 's (as studied in [SF], [DM₂], [J] and [BJ]) in this case of a singleton singular ϕ , and thus to use the counterexample of [BH₂] to show that **the polynomials in S_1 do not characterize any more the approximation order**, leaving perhaps only little hope for a clear and simple characterization of the approximation order for this more general setting.

We use standard multivariate notations. So $\mathbb{Z}_+^s := \{\alpha \in \mathbb{Z}^s : \alpha \geq 0\}$, and for $\alpha \in \mathbb{Z}_+^s$, $(\cdot)^\alpha$ is the power function $x \mapsto x^\alpha := x_1^{\alpha_1} \dots x_s^{\alpha_s}$. D and E are used for the differential resp. shift operators, and so $p(D)$ and $p(E)$ are the differential resp. difference operators obtained by evaluating the polynomial p at D and E ; in particular, $E^\alpha : f \mapsto f(\cdot + \alpha)$.

One concluding remark seems to be in order: the Strang-Fix Conditions, as well as their proofs, strongly emphasize the Fourier analysis approach (Poisson's summation, Parseval formula). Through the years it led to the understanding that the Fourier analysis methods are essential to the characterization of approximation orders and related issues, at least for general uniform meshes. The results here, as well as their proofs, provide an illustration of the alternative techniques: while the Fourier transform approach focuses (implicitly) on the similarity between the semi-discrete operator $\phi \star'$ and the standard convolution operator $\phi \star$ for suitable spaces of polynomials or exponentials (see [BR₃] which elaborates on this point), the alternative approach (as originated in [B₁], with the results in [CJW] and [CD] serving as the motivation) associates $\phi \star'$ with the difference operator

$$T_\phi : f \mapsto \sum_{\alpha \in \mathbb{Z}^s} \phi(\alpha) f(\cdot - \alpha).$$

In this setting, *invariant* subspaces of S_h (under differentiation, $h\mathbb{Z}^s$ -translations, etc.) play a key role. The above difference operators match the action of $\phi \star'$ for more general spaces rather than polynomials or exponentials ([B₁], [R₁]), hence are adequate for the analysis here. Moreover, even for piecewise-polynomials or piecewise-exponentials this approach is sometimes more direct and more efficient than Fourier analysis (cf. e.g., [BAR] and [R₂]).

2. Approximation order for continuous functions

In this section we illustrate the general results by treating first the approximation order for bounded uniformly continuous functions.

Throughout this section, as well as in the next one, we assume that $\{\phi_h\}_h$ is a collection of compactly supported functions which satisfy, for every h , the following three conditions:

$$(2.1) \quad \text{supp } \phi_h \subset [0, hk)^s;$$

$$(2.2) \quad \sum_{\alpha \in h\mathbb{Z}^s} \phi_h(\alpha) = 1;$$

$$(2.3) \quad \|\phi_h\|_\infty \leq c.$$

Here k and c are some constants, which will be used in the sequel in the above meaning without further reference. Note that $\#(h\mathbb{Z}^s \cap \text{supp } \phi_h) \leq k^s$. We also use the notation

$$(2.4) \quad \text{dist}(f, S) := \inf\{\|f - g\|_\infty : g \in S\}.$$

(2.5)Proposition. For every h ,

$$(2.6) \quad \|\phi *_h' 1 - 1\|_\infty \leq 2k^s c \operatorname{dist}(1, S_h).$$

Proof: Let f be an arbitrary element of S_h . Then, for every $\alpha \in \mathbb{R}^s$,

$$(2.7) \quad \|E^{-\alpha} f - 1\|_\infty = \|f - 1\|_\infty.$$

Since $f \in S_h$, we know from [B₁] that $\phi_h *_h' f = \sum_{\alpha \in h\mathbb{Z}^s} \phi_h(\alpha) E^{-\alpha} f$, and hence, by (2.1), (2.2) and (2.7),

$$\|\phi_h *_h' f - 1\|_\infty = \left\| \sum_{\alpha \in h\mathbb{Z}^s} \phi_h(\alpha) (E^{-\alpha} f - 1) \right\|_\infty \leq \sum_{\alpha \in h\mathbb{Z}^s} |\phi_h(\alpha)| \|E^{-\alpha} f - 1\|_\infty \leq k^s c \|f - 1\|_\infty.$$

But this completes the proof, since (2.1) when combined with (2.3), implies that

$$\|\phi_h *_h' (f - 1)\|_\infty \leq k^s c \|f - 1\|_\infty. \quad \spadesuit$$

The above result is now used in the following theorem, which shows that the operators $\{\phi_h *_h'\}_h$ approximate continuous functions to the same order as the best approximations from $\{S_h\}_h$ do.

(2.8) Theorem. Assume that (2.1), (2.2) and (2.3) hold. Then for every function f and every h ,

$$(2.9) \quad \|\phi_h *_h' f - f\|_\infty \leq ck^s (k \omega_f(h) + 2 \operatorname{dist}(1, S_h) \|f\|_\infty),$$

where ω_f is the modulus of continuity of f (in the ∞ -norm). In particular, if $\operatorname{dist}(1, S_h) = o(1)$, then $\operatorname{dist}(f, S_h) = o(1)$ for every bounded uniformly continuous f , and if $\operatorname{dist}(1, S_h) = O(h)$, then $\operatorname{dist}(f, S_h) = O(h)$, for every $f \in C^1(\mathbb{R}^s)$ which is bounded together with its first order derivatives.

Proof: Fix h and $x \in \mathbb{R}^s$. Since, by (2.1), $\phi_h(x - \alpha) = 0$ for all $\alpha \notin x - [0, hk)^s$, then (2.5)Proposition implies that, with $\nu_h(x) := h\mathbb{Z}^s \cap (x - \operatorname{supp} \phi_h)$,

$$(2.10) \quad \left| \sum_{\alpha \in \nu_h(x)} \phi_h(x - \alpha) - 1 \right| \leq 2k^s c \operatorname{dist}(1, S_h),$$

and hence

$$(2.11) \quad \left| \sum_{\alpha \in \nu_h(x)} f(x) \phi_h(x - \alpha) - f(x) \right| \leq 2k^s c \operatorname{dist}(1, S_h) \|f\|_\infty.$$

On the other hand, by (2.3),

$$(2.12) \quad \begin{aligned} & \left| \sum_{\alpha \in \nu_h(x)} f(\alpha) \phi_h(x - \alpha) - \sum_{\alpha \in \nu_h(x)} f(x) \phi_h(x - \alpha) \right| \\ & \leq \sum_{\alpha \in \nu_h(x)} |f(\alpha) - f(x)| |\phi_h(x - \alpha)| \leq ck^{s+1} \omega_f(h). \end{aligned}$$

Summing (2.11) and (2.12) we obtain (2.9), which implies the rest of the theorem. \spadesuit

(2.8)Theorem leads to the following improvement of the case $d = 0$ (and $p = \infty$) in (1.1)Theorem:

(2.13) Corollary. *Let ϕ be a bounded compactly supported function satisfying $\sum_{\alpha \in \mathbb{Z}^s} \phi(\alpha) = 1$. Define $\phi_h := \sigma_{1/h} \phi$, and let S_h be the space spanned by the $h\mathbb{Z}^s$ -translates of ϕ_h . Then the following conditions are equivalent*

- (a) $\text{dist}(f, S_h) = O(h)$, for every $f \in C^1(\mathbb{R}^s)$ which is bounded together with its first order derivatives.
- (b) For every function f

$$\|\phi_h *'_h f - f\|_\infty \leq M \omega_f(h),$$

where M depends only on $\text{diam supp } \phi$ and $\|\phi\|_\infty$.

- (c) $1 \in S_1$.
- (d) $\phi *' 1 = 1$.

Proof:

(d) \implies (c): trivial.

(c) \implies (b): (2.2) is assumed here and (2.1) and (2.3) are trivially satisfied, and therefore (2.8) Theorem can be invoked. (b) then follows from the fact that, since $1 \in S_1$, $1 \in S_h$ for every h , and thus $\text{dist}(1, S_h) = 0$.

(b) \implies (a): trivial.

(a) \implies (d): With $f = 1$, we employ (2.5) Proposition to conclude that $\|\phi_h *'_h 1 - 1\|_\infty \xrightarrow{h \rightarrow 0} 0$.

But since $\|\phi_h *'_h 1 - 1\|_\infty$ is independent of h , it must be 0. ♠

3. Approximation order for smooth functions

In this section we establish necessary and sufficient conditions for higher approximation orders (i.e., higher than $O(h)$) for smooth functions by the spaces $\{S_h\}_h$. Precisely, we look for the maximal d for which $\text{dist}(f, S_h) = O(h^d)$ for every function in the Sobolev space

$$(3.1) \quad W_\infty^d := \{f \in C^d(\mathbb{R}^s) : \|f\|_{\infty, d} := \sum_{|\alpha| \leq d} \|D^\alpha f\|_\infty < \infty\}.$$

Throughout the section we retain assumptions (2.1), (2.2) and (2.3).

The proofs in the previous section were based on the fact that the sequence $\{\phi_h *'_h\}_h$ forms a collection of uniformly bounded local operators which approximate continuous function to the same order as the best approximation. The constant function is a convenient trial function since on the one hand it spans a translation-invariant space while on the other hand it is admissible, being bounded and uniformly continuous. In the general case, though, the operator $\phi_h *'_h$ cannot be expected to approximate well, and will be replaced by an operator of the form $\psi_h *'_h$ for a suitable function $\psi_h \in S_h$. Furthermore, we avoid here polynomials as trial functions, since in general they are neither bounded nor uniformly continuous. Seeking for finite-dimensional spaces H which are translation-invariant on the one hand, and consist of bounded functions on the other hand, we note that such spaces are necessarily of the form

$$(3.2) \quad H = \text{span}\{e_\theta\}_{\theta \in \Theta},$$

where $e_\theta : x \mapsto e^{\theta \cdot x}$, and the **spectrum** Θ of H satisfies

$$(3.3) \quad \Theta \subset i\mathbb{R}^s.$$

The first result here deals with the extension of (2.5) Proposition to exponential spaces H of the above form, and is the crux in the analysis to follow. We use here the notation T_h for the difference operator

$$(3.4) \quad T_h : f \mapsto \sum_{\alpha \in h\mathbb{Z}^s} \phi_h(\alpha) E^{-\alpha} f.$$

(3.5) Theorem. Assume $\{\phi_h\}_h$ satisfy (2.1), (2.2) and (2.3). Let H be a finite-dimensional exponential space satisfying (3.2) and (3.3). Then there exists a sequence $\{V_h\}_h$ of difference operators, supported on $h\mathbb{Z}^s$ (resp.) and satisfying

- (a) V_h inverts $T_{h|_H}$ for all (sufficiently small) h ;
- (b) $\{V_h\}_h$ are uniformly local and uniformly bounded, i.e.,

$$(3.6) \quad \text{supp } V_h \phi_h \subset [0, hk_1)^s, \quad \|V_h \phi_h\|_\infty \leq c_1,$$

for some constants k_1, c_1 .
The functions $\psi_h := V_h \phi_h$ then satisfy

$$(3.7) \quad \|\psi_h *_h' f - f\|_\infty \leq M \text{dist}(f, S_h)$$

for every $f \in H$. The constant M depends only on k, c and $\rho(H) := \max\{\|\theta\|_\infty : e_\theta \in H\}$.

(3.8) Remark. The representation, hence the extension, of $(T_{h|_H})^{-1}$ as difference operators in the approach here is merely a convenient choice. Any uniformly bounded uniformly local extensions will do as well.

Proof: Set Θ for the spectrum of H . With T_h as in (3.4), (2.3) and (2.1) imply that

$$(3.9) \quad \|T_h\| \leq ck,$$

when regarding T_h , say, as an endomorphism of $L_\infty(\mathbb{R}^s)$. We claim that for sufficiently small h , the norms of the restrictions $T_{h|_H}$ are equi-bounded **below** by some positive constant. We use it to define the difference operators $\{V_h\}_h$, which are uniformly bounded (at least for small enough h) and invert $\{T_h\}_h$ on H , (as done in [DR] in the derivation of the approximation order for exponential box splines. The idea, for a polynomial H and $h = 1$, was originated in [CD]). First, we examine the action of T_h on the pure exponentials of H : for an exponential e_θ , $T_h(e_\theta) = e_\theta \sum_{\nu_h(0)} \phi_h(\alpha) e^{-\theta \cdot \alpha}$ (with $\nu_h(0) := h\mathbb{Z}^s \cap [0, hk)^s$), while

$$(3.10) \quad \left| \sum_{\nu_h(0)} \phi_h(\alpha) e^{-\theta \cdot \alpha} \right| \geq \left| \sum_{\nu_h(0)} \phi_h(\alpha) \right| - \left| \sum_{\nu_h(0)} \phi_h(\alpha) (1 - e^{-\theta \cdot \alpha}) \right| \geq 1 - k^s cbh,$$

using (2.1), (2.2) and (2.3). Here b depends on $\rho(H)$ and k but independent of h , which means that (for sufficiently small h and) for all $e_\theta \in H$

$$(3.11) \quad T_h e_\theta = c_{h,\theta} e_\theta, \text{ where } |c_{h,\theta}| \geq 1/2.$$

We now define

$$(3.12) \quad V_h := (I - \prod_{\theta \in \Theta} (I - (1/c_{h,\theta})T_h))/T_h,$$

with I being the identity. Since, for each θ , $I - (1/c_{h,\theta})T_h$ annihilates e_θ , the product term in the definition of V_h annihilates H , and hence, for every h , $(V_h T_h)|_H = I$, which proves (a).

As for the functions $\{\psi_h\}_h$ in the theorem, with the aid of (3.9) and (3.11), we see that $\{V_h\}_h$ is a bounded set of operators, and since each V_h is a linear combination of $\{T_h^j\}_{j=0}^{\#\Theta-1}$, it is a finite difference operator supported on $[0, hr]^s \cap h\mathbb{Z}^s$ (where $r \leq k\#\Theta$), and hence (b) is valid.

To prove (3.7), we now take any $f \in H$ and an arbitrary $f_h \in S_h$. Since

$$(3.13) \quad \|\psi_h *_h' f - f\|_\infty \leq \|\psi_h *_h' f - V_h T_h f_h\|_\infty + \|V_h T_h f_h - f\|_\infty,$$

it is sufficient to estimate each of the right-hand summands in (3.13).

As for the first one, we recall that since $f_h \in S_h$ then $[B_1]$, $T_h f_h = \phi_h *_h' f_h$ and hence $V_h T_h f_h = \psi_h *_h' f_h$. Therefore

$$(3.14) \quad \|\psi_h *_h' f - V_h T_h f_h\|_\infty = \|\psi_h *_h' (f - f_h)\|_\infty \leq k_1^s c_1 \|f - f_h\|_\infty.$$

For the second term, we use the fact that V_h inverts T_h on H , hence

$$(3.15) \quad \|V_h T_h f_h - f\|_\infty = \|V_h T_h (f_h - f)\|_\infty \leq \|V_h T_h\| \|f_h - f\|_\infty.$$

Summing (3.14) and (3.15), and in reliance on the boundedness of $\{V_h T_h\}_h$, we obtain

$$\|\psi_h *_h' f - f\|_\infty \leq M \|f - f_h\|_\infty,$$

which implies (3.7), since f_h was arbitrary. ♠

For later use, we examine in the following corollary the above construction under the scaling assumption.

(3.16) Corollary. *Under the assumption $\phi_h = \sigma_{1/h} \phi_1$, all h , we have*

$$\sigma_h \psi_h \xrightarrow{h \rightarrow 0} ((I - (I - T_1)^{\#\Theta})/T_1) \phi_1,$$

where $\{\psi_h\}_h$ are the functions constructed in (3.5)Theorem. The convergence is uniform.

Proof: Using the notations of (3.5)Theorem we have

$$\sigma_h \psi_h = \sigma_h V_h \sigma_{1/h} \phi_1.$$

Furthermore, the assumption $\phi_h = \sigma_{1/h}\phi_1$ also implies that $\sigma_h T_h \sigma_{1/h} = T_1$, and the claim then follows from the fact that the constants $c_{h,\theta}$ involved in the definition of V_h satisfy $\lim_{h \rightarrow 0} c_{h,\theta} = 1$, $\forall \theta \in \Theta$ ((3.10) bounds this limit above by 1, and a similar argument provides the converse inequality). \spadesuit

The proofs of the main results make use of the “local approximation order”^{*} notion (as do all quasi-interpolation arguments). As a preparation, we define now the **local approximation order** (or the “approximation order at a point”), and discuss some aspects of the local order which are relevant to the context here.

(3.17) Definition. *Let H be a subspace of $C^\infty(\mathbb{R}^s)$. We say that H admits a local approximation order d at $x \in \mathbb{R}^s$ if for every f , which is sufficiently smooth in a neighborhood of x , there exists $g \in H$ such that*

$$(3.18) \quad |(f - g)(x + y)| \leq c_{f,x} \|y\|_\infty^d,$$

for sufficiently small y .

Note that in case H is translation-invariant, the local approximation order of H is independent of x . For an exponential H of the form (3.2), the results of [BR₁] provide a useful characterization of the local approximation order.

(3.19) Result. *Assume H satisfies (3.2). Then the local approximation order d of H is determined by the largest polynomial space π_{d-1} for which the spectrum of H is total.*

We recall that a set Ω is **total** for the function space F , if no $f \in F \setminus 0$ vanishes identically on Ω .

For our purposes, it is also important to have some bound on the approximant g in terms of the Sobolev norm $\|f\|_{\infty,d}$. The proof of Theorem 3.1 of [DR] provides this necessary information, which is specialized to the present context in the following lemma:

(3.20) Result. *Let H be a finite-dimensional exponential space of local approximation order d , and which satisfies (3.2) and (3.3). Then, for every $f \in W_\infty^d$ and $x \in \mathbb{R}^s$, the local approximant $g \in H$ that satisfies (3.18) can be chosen such that*

$$|(f - g)(x + y)| \leq \text{const.} \|f\|_{\infty,d} \|y\|_\infty^d,$$

where const. depends only on H .

(3.21) Theorem. *Suppose that $\{\phi_h\}_h$ satisfy (2.1), (2.2) and (2.3). Let H be a finite-dimensional exponential space that satisfies (3.2) and (3.3) and is of local approximation order d . Let $\{\psi_h\}_h$ be the functions associated with H by (3.5) Theorem. Then, for some constants M_1, M_2 and for every $f \in W_\infty^d$,*

$$(3.22) \quad \|\psi_h *_h' f - f\|_\infty \leq \|f\|_{\infty,d} (M_1 \|h\|_\infty^d + M_2 \max_{e_\theta \in H} \text{dist}(e_\theta, S_h)).$$

^{*} Unfortunately, the terminology “local approximation” is used in spline theory in two different contexts: the “local approximation” mentioned in the introduction in conjunction with [BJ], is thus a different notion.

Proof: Let $f \in W_\infty^d$ and $x \in \mathbb{R}^s$. Then, by (3.5)Theorem and (3.20)Result there exists $g \in H$ for which

$$\begin{aligned} |(\psi_h *'_h f - f)(x)| &\leq |\psi_h *'_h (f - g)(x)| + |(\psi_h *'_h g - g)(x)| \\ &\leq C_1 \|f - g\|_{\infty, x+[0, hr]^s} + C_2 |g(x)| \max_{e_\theta \in H} \text{dist}(e_\theta, S_h) \\ &\leq \|f\|_{\infty, d} (M_1 \|h\|_\infty^d + M_2 \max_{e_\theta \in H} \text{dist}(e_\theta, S_h)), \end{aligned}$$

where in the last inequality, the fact that $g(x) = f(x)$ has been used. ♠

We note that the above theorem is valid (with the same constants) with $\|f\|_{\infty, d}$ replaced by $\|f\|_\infty + \sum_{|\alpha|=d} \|D^\alpha f\|_\infty$.

Combining (3.19)Result with (3.21)Theorem we conclude the following:

(3.23) Corollary. *Suppose that $\{\phi_h\}_h$ satisfy (2.1), (2.2) and (2.3). Then the following conditions are equivalent*

- (a) *For every $f \in W_\infty^d$, $\text{dist}(f, S_h) = O(h^d)$.*
- (b) *For any (every) finite $\Theta \subset i\mathbb{R}^s$ which is total for π_{d-1} , the functions $\{\psi_h\}_h$ constructed in (3.5)Theorem with respect to $H := \text{span}\{e_\theta\}_{\theta \in \Theta}$ satisfy $\|\psi_h *'_h e_\theta - e_\theta\|_\infty = O(h^d)$, for every $\theta \in \Theta$.*

The above corollary gives rise to the following scheme for the determination of the approximation order from the directed spaces $\{S_h\}_h$:

(3.24) Scheme. *Choose*

$$\Theta_1 \subset \Theta_2 \subset \dots \subset i\mathbb{R}^s,$$

such that each Θ_d is total for π_{d-1} and define $H_d := \text{span}\{e_\theta\}_{\theta \in \Theta_d}$. Then for $d = 1, 2, \dots$ do:

Step 1: Check your previous storage of approximation orders, if one of these is d , then stop. d is your approximation order. (Skip this step for $d = 1$).

*Step 2: Construct the quasi-interpolants $\{\psi_h *'_h\}_h$.*

*Step 3: For each $\theta \in \Theta_d \setminus \Theta_{d-1}$, determine the order of $\|\psi_h *'_h e_\theta - e_\theta\|_\infty$. If one of these rates is d , stop. d is then your approximation order. Otherwise store the smallest rate obtained here, increase d by 1 and go to step 1. At this point you already know that the approximation order is at least $d + 1$, and does not exceed the least of the orders stored.*

One way to choose the sets $\{\Theta_d\}_d$ is

$$\Theta_d = \{i\alpha : \alpha \in \mathbb{Z}_+^s, |\alpha| \leq d - 1\}.$$

It should be emphasized that in practice the above results do not necessarily require the identification of the approximation order for constructing good quasi-interpolants. Choosing H of local approximation d , we may use $\{\psi_h\}_h$ of (3.5)Theorem, knowing that the order of the scheme would be $\min\{d, d_0\}$ where d_0 is the (presumably unknown) approximation order.

The last result of this section provides the following improvement of (1.1)Theorem (for $p = \infty$)

(3.25) Theorem. Let ϕ be a compactly supported bounded function satisfying $\sum_{\alpha \in \mathbb{Z}^s} \phi(\alpha) \neq 0$. Define $\phi_h := \sigma_{1/h} \phi$, and $S_h := \text{span}\{E^\alpha \phi_h\}_{\alpha \in h\mathbb{Z}^s}$. Then the following conditions are equivalent:
(a) For every function $f \in W_\infty^d$

$$(3.26) \quad \text{dist}(f, S_h) = O(h^d).$$

(b) $\pi_{d-1} \subset S_1$.

Proof: The implication (b) \implies (a) is well-known (and by now standard). For the converse, we choose Θ to be a finite subset of $i\mathbb{R}^s$ which is total for π_{d-1} and define $H := \text{span}\{e_\theta\}_{\theta \in \Theta}$. Then, since $H \subset W_\infty^d$, $\text{dist}(f, S_h) = O(h^d)$ for every $f \in H$ and hence (3.21)Theorem (when combined with (3.19)Result) implies that the $\{\psi_h\}_h$ constructed in (3.5)Theorem (with respect to the present H) satisfy

$$\|\psi_h *'_h f - f\|_\infty = O(h^d),$$

for every $f \in W_\infty^d$. Rescaling each ψ_h back to the original mesh size we obtain a set of function $\{g_h := \sigma_h \psi_h\}_h \subset S_1$, which, by (3.16)Corollary, converges uniformly to some $g \in S_1$.

Now, let $\gamma \in \mathbb{Z}_+^s$, and choose $f \in W_\infty^d$ which coincides with a homogeneous polynomial p of degree k in a neighbourhood of the origin. Then, for a fixed $x \in \mathbb{R}^s$ and sufficiently small h ,

$$\begin{aligned} h^k |(g_h *'_h p - p)(x)| &= h^k |(\psi_h *'_h \sigma_{1/h} p)(hx) - p(x)| \\ &= |(\psi_h *'_h p - p)(hx)| \\ &= |(\psi_h *'_h f - f)(hx)| \\ &\leq \|\psi_h *'_h f - f\|_\infty = O(h^d). \end{aligned}$$

Thus, in case $k \leq d-1$, we obtain

$$(g *'_h p - p)(x) = \lim_{h \rightarrow 0} (g_h *'_h p - p)(x) = 0.$$

We conclude that $g *'_h p = p$ for every homogeneous $p \in \pi_{d-1}$, hence for every $p \in \pi_{d-1}$, and since $g \in S_1$, $\pi_{d-1} \subset S_1$ as well. \spadesuit

4. Singularity and the case of several compactly supported functions

Our primary aim in this section is to obtain results concerning the approximation order in case $\phi := \phi_1$ satisfies the singularity condition

$$(4.1) \quad \sum_{\alpha \in \mathbb{Z}^s} \phi(\alpha) = 0,$$

and the refined spaces $\{S_h\}$ are obtained by scaling. The questions we focus on are whether good approximation rates are possible at all in such a situation (answer: yes), and whether the approximation order is determined by the polynomials in $S(\phi) := S_1(\phi)$ in the same way as in the regular case (answer: in general, no).

Some of the methods employed here are aimed to connect the case of a single ϕ to the case of many ϕ 's. In doing so, we obtain several results on the latter case which are of independent interest.

Throughout this section ϕ is a compactly supported bounded function, $S(\phi)$ is the space spanned by the integer translates of ϕ , and the spaces $\{S_h\}_h := \{S_h(\phi)\}_h$ are obtained by scaling $S(\phi)$, i.e., $S_h = \sigma_{1/h} S(\phi)$.

We first show that (4.1) dashes any hope for a positive approximation order, unless the stronger assumption

$$(4.2) \quad \phi *' 1 = 0,$$

is made:

(4.3) Proposition. *Assume that (4.1) holds, yet $\phi *' 1 \neq 0$. Then $\text{dist}(1, S_h) \neq O(h)$.*

Proof: Since 1 is translation-invariant, then $\text{dist}(1, E^x S_h) = \text{dist}(1, S_h)$, for every h and every real translation E^x of S_h . On the other hand, since $\phi *' 1 \neq 0$, one can find a translate $\psi := E^x \phi$ which satisfies the regularity condition $\sum_{\alpha \in \mathbb{Z}^s} \psi(\alpha) \neq 0$. Observing that $S_h(\psi) = E^{hx} S_h(\phi)$, we conclude that $\text{dist}(1, S_h(\phi)) = \text{dist}(1, S_h(\psi))$ for all h . As for ψ , this function is regular, hence (2.13)Corollary implies that $\text{dist}(1, S_h(\psi)) \neq O(h)$, unless $\psi *' 1 = \text{const.} \neq 0$. Yet, the latter implies $\phi *' 1 = \text{const.} \neq 0$, which is impossible, since by (4.1) $(\phi *' 1)|_{\mathbb{Z}^s} = 0$. ♠

To this end, we assume therefore that (4.2) holds and mention that already the authors of [SF] indicated that (4.2) does not contradict good approximation orders (as a matter of fact, [SF] identifies the “bad case” as being $\hat{\phi}(0) = 0$, a condition which is implied by (4.2)). Indeed, let ϕ be a **regular** compactly supported function, and p a linear polynomial, $p(1, \dots, 1) = 0$. Defining $\psi := p(E)\phi$, we see that ψ is singular, yet $S(\phi) = S(\psi)$, hence the approximation rates from the scales of $S(\psi)$ can be arbitrarily high, as the approximation rates associated with $S(\phi)$ can.

In general, not every compactly supported singular spline is obtained by differencing a regular one (the results in the sequel will prove this indirectly). However, in the **univariate** case, every compactly supported function that satisfies (4.2) is obtained by differencing another function. This leads to the following

(4.4) Proposition. *For a univariate ϕ , (3.25)Theorem holds even without the assumption $\sum_{\alpha \in \mathbb{Z}} \phi(\alpha) \neq 0$.*

Proof: There is nothing to prove unless (4.1) holds. Moreover, if (4.2) does not hold, then (4.3)Proposition shows that both (a) and (b) of (3.25)Theorem are false for every positive d . We can therefore assume that (4.2) is valid. Now, choose k such that $\text{supp } \phi \subset [0, k)$. Invoking Proposition 3.2 of [R₂], we conclude that $\phi = (E - 1)\psi$ for some function ψ supported in $[0, k - 1)$. Since $S(\phi) = S(\psi)$, we may replace ϕ by ψ for determining the approximation order. If ψ satisfies $\psi *' 1 = 0$, the process can be repeated. After at most $k - 1$ steps, we obtain a function τ with $S(\tau) = S(\phi)$, which necessarily satisfies $\tau *' 1 \neq 0$, hence we are reduced to the former case. ♠

We are aiming now at proving the following

(4.5) **Claim.** *There exists a singular compactly supported function ϕ for which the implication (b) \implies (a) of (3.25)Theorem is invalid.*

By the results so far, such ϕ cannot be univariate, and must satisfy (4.2). We will not construct this ϕ explicitly. Instead we draw here a link between our setting and the more involved case of several compactly supported functions, and make use of (negative) results known in the latter case.

The setting is as follows: $\Phi := \{\phi_j\}_{j=1}^n$ is a set of compactly supported functions. $S(\Phi)$ is the space spanned by their integer translates. An **exponential space** here is a space H which satisfies

$$(4.6) \quad H \subset \sum_{\theta \in \Theta} e_\theta \pi,$$

for some finite $\Theta \subset C^s$. The minimal possible Θ in (4.6) is the **spectrum** of H . An **exponential** is an element of an exponential space. For a compactly supported ϕ , we define $H(\phi)$ as the space of all exponentials in $S(\phi)$. $H(\phi)$ is of importance in multivariate splines, since its local approximation properties may lead to lower bounds on the approximation order attained by appropriate refined versions of $S(\phi)$ ([DR], [BR₃]). Here, we always assume that $H(\phi)$ is **D-invariant**, i.e., closed under differentiations; this implies that $H(\phi)$ can be decomposed into

$$(4.7) \quad H(\phi) = \bigoplus_{\theta \in \Theta} e_\theta P_{\phi, \theta},$$

where $P_{\phi, \theta}$ are finite-dimensional D -invariant polynomial spaces. Our result is as follows

(4.8) **Theorem.** *Let Φ be as above and assume that $H(\phi_j)$ is D -invariant for $j = 1, \dots, n$. Then, for $s > 1$, there exists a compactly supported $\psi \in S(\Phi)$ that satisfies*

$$(4.9) \quad H(\phi_j) \subset H(\psi), \quad j = 1, \dots, n.$$

First, we show how the above theorem is connected to (4.5)Claim. In the above terminology, a polynomial space is a special case of an exponential space. For a polynomial space, [B₁] proves that $S(\phi) \cap \pi$ is always D -invariant, and so (4.8)Theorem shows in particular that there exists a compactly supported $\psi \in S(\Phi)$, which satisfies

$$(4.10) \quad S(\phi_j) \cap \pi \subset S(\psi), \quad j = 1, \dots, n.$$

Now, the example given in [BH₂] provides an instance of a set $\Phi = \{\phi_1, \phi_2\}$ of two bivariate box splines such that $\pi_3 \subset S(\Phi)$, yet the corresponding approximation order from $\{\sigma_{1/h} S(\Phi)\}_h$ is only 3, (and not 4 as might have been expected). By (4.8)Theorem, there exists $\psi \in S(\Phi)$ satisfying

$$\pi_3 \subset S(\psi),$$

while the approximation order from $S_h(\psi)$ is at most 3, since $S(\psi)$ is a subspace of $S(\Phi)$.

In the rest of this section we prove (4.8)Theorem, and discuss other problems initiated by the theorem and related to the case of several ϕ 's. The discussion of approximation orders for $\{S_h\}_h$ ends therefore at this point. Note that we were unable to comment on the validity of the implication (a) \implies (b) in (3.25)Theorem in the case (4.2) is assumed.

We turn now to the proof of (4.8)Theorem. This proof is based on the following result from [BR₂]:

(4.11) **Result.** Let $e_\theta P$, $P \subset \pi$ be a D -invariant subspace of $S(\phi)$, ϕ being compactly supported. Then, there exists a finite-dimensional D -invariant $Q \subset \pi$ such that $e_\theta Q$ is mapped by $\phi *'$ onto $e_\theta P$.

The theorem now follows by a repeated application of the following lemma:

(4.12) **Lemma.** Let ϕ_1, ϕ_2 be two compactly supported functions defined on \mathbb{R}^s where $s > 1$. Let H_1, H_2 be two D -invariant exponential spaces. If

$$(4.13) \quad H_j \subset S(\phi_j), \quad j = 1, 2,$$

then there exists a compactly supported $\psi \in S(\{\phi_1, \phi_2\})$ such that

$$(4.14) \quad H_j \subset S(\psi), \quad j = 1, 2.$$

Proof: Set Θ_j for the spectrum of each H_j . Since each H_j is D -invariant, it is a direct sum of spaces of the form $e_\theta P_\theta$, where $\theta \in \Theta_j$ and P_θ is a D -invariant polynomial space. Hence, by (4.11) Result, there exist finite-dimensional exponential F_j with spectrum Θ_j , $j = 1, 2$, such that

$$\phi_j *' F_j = H_j.$$

Define $v^1 := (1, 0, \dots, 0)$, $v^2 := (0, 1, 0, \dots, 0)$. Let T_j , $j = 1, 2$, be two non-trivial finite difference operators such that, for $j = 1, 2$,

- (a) T_j is supported on $v^j \mathbb{Z}$;
- (b) $T_1 F_2 = T_2 F_1 = 0$.

Each T_j is a finite difference operator, and therefore for sufficiently high k , $T_j(e_\theta(\cdot)^{kv^j}) \neq 0$, for all $\theta \in \Theta_1 \cup \Theta_2$, which implies that $T_j U_j$ is 1-1 on $e_\theta \pi$ for each $\theta \in \Theta_1 \cup \Theta_2$, where

$$U_j : f \mapsto (\cdot)^{kv^j} f, \quad j = 1, 2.$$

Now, let V_j be a finite difference operator which inverts $(T_j U_j)|_{F_j}$ from the left, and define $G_j := V_j U_j F_j$. Note that T_1 commutes with V_1, V_2 and U_2 , hence

$$T_1 G_2 = V_2 U_2 T_1 F_2 = 0, \quad T_1 G_1 = T_1 V_1 U_1 F_1 = V_1 T_1 U_1 F_1 = F_1,$$

and similarly $T_2 G_1 = 0$, $T_2 G_2 = F_2$. Define

$$\psi = \sum_{j=1}^2 T_j \phi_j.$$

Then, by the above,

$$\psi *' G_1 = \phi_1 *' (T_1 G_1) + \phi_2 *' (T_2 G_1) = \phi_1 *' F_1 = H_1,$$

and similarly $\psi *' G_2 = H_2$. ♠

(4.15) **Corollary.** Assume $s > 1$. Then every finite-dimensional D -invariant exponential space is $H(\phi)$ for some compactly supported ϕ .

Proof: From [R₁; Thm. 1.1] we know that every finite-dimensional D -invariant exponential subspace of $e_\theta\pi$ (with $\theta \in \mathbb{C}^s$) is $H(\phi)$ for some compactly supported ϕ . (4.8) Theorem then completes the proof, since every finite-dimensional D -invariant exponential space is the sum of D -invariant spaces of the form $e_\theta P \subset e_\theta\pi$. \spadesuit

The fact that the univariate case was not covered here is essential and is not related to the technique used (cf. Proposition 4.6 in [R₂], which shows that not every univariate D -invariant exponential space is $H(\phi)$ for some ϕ). For completeness, we record the following result, which under some “regularity” assumption (where regularity is now in the sense of [R₂]) enables us to obtain more, and in particular to include the univariate case.

(4.16) **Theorem.** Let $\Phi = \{\phi_1, \dots, \phi_n\}$ be a collection of n compactly supported functions. For $j = 1, \dots, n$, let H_j be a D -invariant exponential subspace of $S(\phi_j)$ with spectrum Θ_j , such that, for $j \neq k$,

$$(4.17) \quad (\Theta_j - \Theta_k) \cap 2\pi i\mathbb{Z}^s = \emptyset.$$

Then there exists a compactly supported $\psi \in S(\Phi)$ with the properties

- (a) $H_j \subset S(\psi), \forall j$;
- (b) $\psi *' e_\theta \neq 0$, in case $\theta \in \Theta_j$ and $\phi_j *' e_\theta \neq 0$.

Proof: As in the previous proof, we first apply (4.11) Result, to find exponential spaces $\{F_j\}_j$ with corresponding spectra $\{\Theta_j\}_j$, such that $\phi_j *' F_j = H_j, j = 1, \dots, n$. The assumption (4.17) then implies, [BAR], [BR₃], the existence of difference operators $\{T_j\}_j$ supported on \mathbb{Z}^s such that

$$(4.18) \quad T_j F_k = 0, \text{ for } j \neq k, \quad \text{and } T_j \text{ is } 1 - 1 \text{ on } \sum_{\theta \in \Theta_j} e_\theta \pi.$$

Defining $\psi = \sum_j T_j \phi_j$, we obtain

$$(4.19) \quad \psi *' F_k = \sum_j \phi_j *' T_j(F_k) = T_k(\phi *' F_k) = T_k H_k.$$

Since H_j is D -invariant, hence translation-invariant, $T_k H_k \subset H_k$. Yet, T_k is 1-1 on $\sum_{\theta \in \Theta_k} e_\theta \pi$, hence is also 1-1 on its subspace H_k . We conclude that $T_k H_k = H_k$, and (a) follows.

For (b), we first note that whenever $j \neq k$, $T_j H_k = \phi_k *' (T_j F_k) = 0$, by (4.18), and hence if $e_\theta \in H_j$ for some j , then, as in (4.19), $\psi *' e_\theta = T_j(\phi_j *' e_\theta)$. Moreover, since e_θ spans a translation-invariant space, [B₁] shows that the assumption $e_\theta \in S(\phi_j)$ implies that $\phi_j *' e_\theta = c e_\theta$, for some constant c , which is non-zero by the assumptions here. Also, T_j is injective on H_j , hence $T_j e_\theta = c_1 e_\theta$ for some non-zero c_1 . Combining all of these we see that

$$\psi *' e_\theta = c c_1 (\phi_j *' e_\theta), \quad c c_1 \neq 0,$$

and (b) follows. ♠

Corollary 2.1 of [R₁] shows that the above theorem is sharp in following sense: whenever $e_\theta, e_\vartheta \in S(\psi)$ and $\theta - \vartheta \in 2\pi i\mathbb{Z}^s \setminus 0$, $\psi *' e_\theta = \psi *' e_\vartheta = 0$.

References

- [B₁] C. de Boor, The polynomials in the linear span of integer translates of a compactly supported function, *Constructive Approximation* **3** (1987), 199–208.
- [B₂] C. de Boor, Quasiinterpolants and approximation power of multivariate splines. In *Computation of curves and surfaces*, M. Gasca and C. A. Micchelli eds., Dordrecht, Netherlands: Kluwer Academic Publishers, (1990), xxx-xxx.
- [BAR] A. Ben-Artzi and A. Ron, Translates of exponential box splines and their related spaces, *Trans. Amer. Math. Soc.* **309** (1988), 683–710.
- [BH₁] C. de Boor and K. Höllig, B-splines from parallelepipeds, *J. d'Anal. Math.* **42** (1982/3), 99–115.
- [BH₂] C. de Boor and K. Höllig, Approximation order from bivariate C^1 -cubics: a counterexample, *Proc. Amer. Math. Soc.* **87** (1983), 649–655.
- [BJ] C. de Boor and R.Q. Jia, Controlled approximation and a characterization of the local approximation order. *Proc. Amer. Math. Soc.* **95** (1985), 547–553.
- [BR₁] C. de Boor and A. Ron, On multivariate polynomial interpolation, CMS TSR 89-17, University of Wisconsin-Madison, November 1988, *Constructive Approximation*, to appear.
- [BR₂] C. de Boor and A. Ron, Polynomial ideals and multivariate splines, CS TR #849, University of Wisconsin-Madison, June 1989, in *Multivariate approximation Theory V*, W. Schemp & K. Zeller eds., Birkhäuser, Basel (1990), xxx-xxx..
- [BR₃] C. de Boor and A. Ron, The exponentials in the span of the integer translates of a compactly supported function, CS TR 887, University of Wisconsin-Madison, November 1989.
- [CD] C. K. Chui and H. Diamond, A natural formulation of quasi-interpolation by multivariate splines, *Proc. Amer. Math. Soc.* **99** (1987), 643–646.
- [CJW] C.K. Chui and K. Jetter, J. Ward, Cardinal interpolation by multivariate splines, *Math. Comp.* **48** (1987), 711–724.
- [DM₁] W. Dahmen and C. A. Micchelli, Translates of multivariate splines. *Linear Algebra and Appl.* **52/3** (1983), 217–234.
- [DM₂] W. Dahmen and C.A. Micchelli, On the approximation order from certain multivariate spline spaces. *J. Austral. Math. Soc. Ser B* **26** (1984) 233–246.
- [DR] N. Dyn and A. Ron, Local Approximation by certain spaces of multivariate exponential-polynomials, approximation order of exponential box splines and related interpolation problems, CAT Rep. 160, Texas A&M Univ., College Station, TX 77843, Jan. 1988. *Trans. Amer. Math. Soc.* **316** (1989), xxx-xxx.
- [J] R.-q. Jia, A counterexample to a result concerning controlled approximation, *Proc. Amer. Math. Soc.* **97** (1986), 647–654.

- [R₁] A. Ron, Relations between the support of a compactly supported function and the exponential-polynomials spanned by its integer translates, *Constructive Approximation*, to appear.
- [R₂] A. Ron, Factorization theorems for univariate splines on regular grids, CS TR #829, University of Wisconsin-Madison, February 1989, *Israel J. Math.*, to appear.
- [SF] G. Strang and G. Fix, A Fourier analysis of the finite element variational method. C.I.M.E. II Cilo 1971, in *Constructive Aspects of Functional Analysis* ed. G. Geymonet 1973, 793-840.