

IDEAL INTERPOLATION: MOURRAIN'S CONDITION VS D -INVARIANCE

C. DE BOOR

POB 1076

Eastsound WA 98245 USA

E-mail: deboor@cs.wisc.edu

Abstract. Mourrain [Mo] characterizes those linear projectors on a finite-dimensional polynomial space that can be extended to an ideal projector, i.e., a projector on polynomials whose kernel is an ideal. This is important in the construction of normal form algorithms for a polynomial ideal. Mourrain's characterization requires the polynomial space to be 'connected to 1', a condition that is implied by D -invariance in case the polynomial space is spanned by monomials. We give examples to show that, for more general polynomial spaces, D -invariance and being 'connected at 1' are unrelated, and that Mourrain's characterization need not hold when his condition is replaced by D -invariance.

By definition (see [Bi]), **ideal interpolation** is provided by a linear projector whose kernel is an ideal in the ring Π of polynomials (in d real ($\mathbb{F} = \mathbb{R}$) or complex ($\mathbb{F} = \mathbb{C}$) variables). The standard example is Lagrange interpolation; the most general example has been called 'Hermite interpolation' (in [M] and [Bo]) since that is what it reduces to in the univariate case.

Ideal projectors also occur in computer algebra, as the maps that associate a polynomial with its *normal form* with respect to an ideal; see, e.g., [CLO]. It is in this latter context that Mourrain [Mo] poses and solves the following problem. *Among all linear projectors N on*

$$\Pi_1(F) := \sum_{j=0}^d ()_j F$$

with range the linear space F , characterize those that are the restriction to $\Pi_1(F)$ of an ideal projector with range F . Here,

$$()_j := ()^{\varepsilon_j}, \quad \varepsilon_j := (\delta_{jk} : k = 1:d), \quad j = 0:d,$$

2000 *Mathematics Subject Classification*: Primary 41A05, 41A10, 41A63; Secondary 13P10.
 The paper is in final form and no version of it will be published elsewhere.

with

$$(\cdot)^\alpha : \mathbb{F}^d \rightarrow \mathbb{F} : x \mapsto x^\alpha := \prod_{j=1}^d x(j)^{\alpha(j)}$$

a handy if nonstandard notation for the **monomial with exponent** α , with

$$\alpha \in \mathbb{Z}_+^d := \{\alpha \in \mathbb{Z}^d : \alpha(j) \geq 0, j = 1:d\}.$$

I also use the corresponding notation

$$D_j$$

for the derivative with respect to the j th argument, and

$$D^\alpha := \prod_{j=1}^d D_j^{\alpha(j)}, \quad \alpha \in \mathbb{Z}_+^d.$$

To state Mourrain's result, I also need the following, standard, notations. The (total) **degree** of the polynomial $p \neq 0$ is the nonnegative integer

$$\deg p := \max\{|\alpha| : \widehat{p}(\alpha) \neq 0\},$$

with

$$p =: \sum_{\alpha} (\cdot)^\alpha \widehat{p}(\alpha),$$

and

$$|\alpha| := \sum_j \alpha(j),$$

while

$$\Pi_{<n} := \{p \in \Pi : \deg p < n\}.$$

THEOREM 1 ([Mo]). *Let F be a finite-dimensional linear subspace of Π satisfying **Mourrain's condition**:*

$$(2) \quad f \in F \implies f \in \Pi_1(F \cap \Pi_{<\deg f}),$$

and let N be a linear projector on $\Pi_1(F)$ with range F . Then, the following are equivalent:

- (a) *N is the restriction to $\Pi_1(F)$ of an ideal projector with range F .*
- (b) *The linear maps $M_j : F \rightarrow F : f \mapsto N((\cdot)_j f)$, $j = 1:d$, commute.*

For a second proof of this theorem and some unexpected use of it in the setting of ideal interpolation, see [Bo].

Mourrain's condition (2) implies that, if F contains an element of degree k , it must also contain an element of degree $k - 1$. In particular, if F is nontrivial, then it must contain a constant polynomial. This explains why Mourrain [Mo] calls a linear subspace satisfying his condition **connected to 1**. Since the same argument can be made in case F is **D -invariant**, i.e., closed under differentiation, this raises the question what connection if any there might be between these two properties.

In particular, for the special case $d = 1$, if F is a linear subspace of dimension n and either satisfying Mourrain's condition or being D -invariant, then, necessarily, $F = \Pi_{<n}$.

More generally, if F is an n -dimensional subspace in the subring generated by the linear polynomial

$$\langle \cdot, y \rangle : \mathbb{F}^d \rightarrow \mathbb{F} : x \mapsto \langle x, y \rangle := \sum_{j=1}^d x(j)y(j)$$

for some $y \neq 0$, then, either way,

$$F = \text{ran}[\langle \cdot, y \rangle^{j-1} : j = 1:n] := \left\{ \sum_{j=1}^n \langle \cdot, y \rangle^{j-1} a(j) : a \in \mathbb{F}^n \right\}.$$

As a next example, assume that F is a **monomial** space (meaning that it is spanned by monomials). If such F is D -invariant, then, with each $()^\alpha$ for which $\alpha - \varepsilon_j \in \mathbb{Z}_+^d$, it also contains $()^{\alpha - \varepsilon_j}$ and therefore evidently satisfies Mourrain's condition.

Slightly more generally, assume that F is **dilation-invariant**, meaning that it contains $f(h \cdot)$ for every $h > 0$ if it contains f or, equivalently, F is spanned by homogeneous polynomials. Then every $f \in F$ is of the form

$$f =: f_{\uparrow} + f_{< \deg f},$$

with f_{\uparrow} the **leading** term of f , i.e., the unique homogeneous polynomial for which

$$\deg(f - f_{\uparrow}) < \deg f,$$

hence in F by dilation-invariance, therefore also

$$f_{< \deg f} \in F_{< \deg f} := F \cap \Pi_{< \deg f},$$

while, by the homogeneity of f_{\uparrow} ,

$$\sum_{j=1}^d ()_j D_j(f_{\uparrow}) = (\deg f) f_{\uparrow}$$

(this is **Euler's theorem for homogeneous functions**; see, e.g., [Enc: p281] which gives the reference [E: §225 on p154]). If now F is also D -invariant, then $D_j(f_{\uparrow}) \in F_{< \deg f}$, hence, altogether,

$$f \in \Pi_1(F_{< \deg f}), \quad f \in F.$$

In other words, *if a dilation-invariant finite-dimensional subspace F of Π is D -invariant, then it also satisfies Mourrain's condition.*

On the other hand, the linear space

$$\text{ran}[(\cdot)^0, (\cdot)^{1,0}, (\cdot)^{1,1}] = \{(\cdot)^0 a + (\cdot)^{1,0} b + (\cdot)^{1,1} c : a, b, c \in \mathbb{F}\}$$

fails to be D -invariant even though it satisfies Mourrain's condition and is monomial, hence dilation-invariant.

The final example, of a space that is D -invariant but does not satisfy Mourrain's condition, is slightly more complicated. In its discussion, I find it convenient to refer to

$$\text{supp } \hat{p}$$

as the **'support'** of the polynomial $p = \sum_{\alpha} ()^\alpha \hat{p}(\alpha)$, with the quotation marks indicating that it is not actually the support of p but, rather, the support of its coefficient sequence,

\widehat{p} . The example is provided by the D -invariant space F generated by the polynomial

$$p = ()^{1,7} + ()^{3,3} + ()^{5,0},$$

hence the ‘support’ of p is

$$\text{supp } \widehat{p} = \{(1, 7), (3, 3), (5, 0)\}$$

(see (4) below). Here are a first few elements of F :

$$D_1 p = ()^{0,7} + 3()^{2,3} + 5()^{4,0}, \quad D_2 p = 7()^{1,6} + 3()^{3,2},$$

hence

$$D_1 D_2 p = 7()^{0,6} + 9()^{2,2}, \quad D_2^2 p = 42()^{1,5} + 6()^{3,1},$$

also

$$D_1^2 p = 6()^{1,3} + 20()^{3,0}, \quad D_1 D_2^2 p = 42()^{0,5} + 18()^{2,1},$$

etc. This shows (see (4) below) that any $q \in \Pi_1(F_{<\text{deg } p})$ having some ‘support’ in $\text{supp } \widehat{p}$ is necessarily a weighted sum of $({}_1 D_1 p)$ and $({}_2 D_2 p)$ (and, perhaps, others not having any ‘support’ in $\text{supp } \widehat{p}$), yet $(p, ({}_1 D_1 p), ({}_2 D_2 p))$ is linearly independent ‘on’ $\text{supp } \widehat{p}$, as the matrix

$$\begin{bmatrix} 1 & 1 & 7 \\ 1 & 3 & 3 \\ 1 & 5 & 0 \end{bmatrix}$$

(of their coefficients indexed by $\alpha \in \text{supp } \widehat{p}$) is evidently 1-1. Consequently,

$$p \notin \Pi_1(F_{<\text{deg } p}),$$

i.e., this F does not satisfy Mourrain’s condition.

This space also provides the proof that, in Theorem 1, one may not, in general, replace Mourrain’s condition by D -invariance.

PROPOSITION 3. *Let F be the D -invariant space spanned by*

$$p = ()^{1,7} + ()^{3,3} + ()^{5,0}.$$

Then there exists a linear projector, N , on $\Pi_1(F)$ with range F for which (b) but not (a) of Theorem 1 is satisfied.

Proof. For $\alpha, \beta \in \mathbb{Z}_+^d$, set

$$[\alpha \dots \beta] := \{\gamma \in \mathbb{Z}_+^d : \alpha \leq \gamma \leq \beta\},$$

with

$$\alpha \leq \gamma := \alpha(j) \leq \gamma(j), \quad j = 1:d.$$

With this, we determine a basis for F as follows.

Since $D^{0,4} p$ is a positive scalar multiple of $({}^1 1, 3)$, we know, by the D -invariance of F , that

$$\{({}^\zeta : \zeta \in [(0, 0) \dots (1, 3)]\} \subset F.$$

This implies, considering $D^{2,0} p$, that $({}^3 0)$, hence also $({}^2 0)$, is in F . Hence, altogether,

$$F = \Pi_{\Xi_0} \oplus \text{ran}[D^\alpha p : \alpha \in [(0, 0) \dots (1, 3)]],$$

Then, none of these is in F , and, among them, each b_ζ is the only one having some ‘support’ at ζ , hence they form a linearly independent sequence. Therefore, each such b_ζ is in b_Z .

The remaining candidates for membership in b_Z require a more detailed analysis. We start from the ‘top’, showing also along the way that (b) of Theorem 1 holds for this F and N by verifying that

$$(5) \quad M_1 M_2 = M_2 M_1 \quad \text{on } b_\xi$$

for every $\xi \in \Xi$.

$\xi = (1, 7)$: As already pointed out, both $()_1 b_{1,7}$ and $()_2 b_{1,7}$ are in b_Z , hence (5) holds trivially for $\xi = (1, 7)$.

$\xi = (0, 7), (1, 6)$: Both $()_1 b_{0,7} = ()^{1,7} + 3()^{3,3} + 5()^{5,0}$ and $()_2 b_{1,6} = 7()^{1,7} + 3()^{3,3}$ have their ‘support’ in that of $p = b_{1,7} = ()^{1,7} + ()^{3,3} + ()^{5,0}$, while, as pointed out and used earlier, the three are independent. Hence $()_1 b_{0,7}, ()_2 b_{1,6} \in b_Z$, while we already pointed out that $()_2 b_{0,7}, ()_1 b_{1,6} \in b_Z$, therefore (5) holds trivially.

$\xi = (0, 6), (1, 5)$: Both $()_1 b_{0,6} = 7()^{1,6} + 9()^{3,2}$ and $()_2 b_{1,5} = 42()^{1,6} + 6()^{3,2}$ have their ‘support’ in that of $b_{1,6} = 7()^{1,6} + 3()^{3,2}$, but neither is a scalar multiple of $b_{1,6}$. Hence, one is in b_Z and the other is not. Which is which depends on the ordering of the columns of $[b_\Xi, ()_1 b_\Xi, ()_2 b_\Xi]$. Assume the ordering such that $()_2 b_{1,5} \in b_Z$. Then, since we already know that $()_1 b_{1,5} \in b_Z$, (5) holds trivially for $\xi = (1, 5)$. Further, $()_1 b_{0,6} = 4b_{1,6} - (1/2)()_2 b_{1,5}$, hence $M_1 b_{0,6} = 4b_{1,6}$, while we already know that $()_2 b_{1,6} \in b_Z$ therefore, $M_2 M_1 b_{0,6} = 0$. On the other hand, $()_2 b_{0,6} = 7()^{0,7} + 3()^{3,3}$ has its ‘support’ in that of $b_{0,7} = ()^{0,7} + 3()^{3,3} + 5()^{4,0}$ but is not a scalar multiple of it, hence is in b_Z , and therefore already $M_2 b_{0,6} = 0$. Thus, (5) also holds for $\xi = (0, 6)$.

$\xi = (0, 5), (1, 4)$: Both $()_1 b_{0,5} = 42()^{1,5} + 18()^{3,1}$ and $()_2 b_{1,4} = 210()^{1,5} + 6()^{3,1}$ have their ‘support’ in that of $b_{1,5} = 42()^{1,5} + 6()^{3,1}$ but $()^{3,1} = b_{3,1}$ was already identified as an element of b_Z , hence neither $()_1 b_{0,5}$ nor $()_2 b_{1,4}$ is in b_Z . But, since $()^{3,1} \in b_Z$, and so $b_{1,5} = N b_{1,5} = N(42()^{1,5})$, we have $M_1 b_{0,5} = b_{1,5}$ and $M_2 b_{1,4} = 5b_{1,5}$. Since we already know that $()_1 b_{1,5} \in b_Z$, it follows that $M_1 M_2 b_{1,4} = 0$ while we already know that $()_1 b_{1,4} \in b_Z$, hence already $M_1 b_{1,4} = 0$. Therefore, (5) holds for $\xi = (1, 4)$. Further, we already know that $()_2 b_{1,5} \in b_Z$, hence $M_2 M_1 b_{0,5} = 0$, while $()_2 b_{0,5} = 42()^{0,6} + 18()^{2,2}$ has the same ‘support’ as $b_{0,6} = 7()^{0,6} + 9()^{2,2}$ but is not a scalar multiple of it, hence is in b_Z and, therefore, already $M_2 b_{0,5} = 0$, showing that (5) holds for $\xi = (0, 5)$.

$\xi = (0, 4)$: $()_2 b_{0,4} = 210()^{0,5} + 18()^{2,1} = 5b_{0,5} - 72b_{2,1}$, with $b_{2,1} \in b_Z$, hence $()_2 b_{0,4}$ is not in b_Z and $M_2 b_{0,4} = 5b_{0,5}$, therefore $M_1 M_2 b_{0,4} = 5M_1 b_{0,5} = 5b_{1,5}$, the last equation from the preceding paragraph. On the other hand, $()_1 b_{0,4} = 210()^{1,4} + 18()^{3,0} = b_{1,4} + 12b_{3,0}$, with both $b_{1,4}$ and $b_{3,0}$ in F , hence $()_1 b_{0,4}$ is not in b_Z , and $M_1 b_{0,4} = b_{1,4} + 12b_{3,0}$, therefore, since $()_2 b_{3,0} = b_{3,1} \in b_Z$, $M_2 M_1 b_{0,4} = M_2 b_{1,4} = 5b_{1,5}$, the last equation from the preceding paragraph. Thus, (5) holds for $\xi = (0, 4)$.

$\xi = (1, 3)$: We already know that $()_1 b_{1,3} = b_{2,3} \in b_Z$ and therefore already $M_1 b_{1,3} = 0$, while $()_2 b_{1,3} = ()^{1,4} = (b_{1,4} - 6b_{3,0})/210 \in F$, therefore $210M_1 M_2 b_{1,3} = M_1 b_{1,4} = 0$, thus (5) holds for $\xi = (1, 3)$.

For the remaining $\xi \in \Xi$, each b_ξ is a monomial, hence $()_j b_\xi$ is again a monomial, and

either in F or not and, if not, then its exponent is in

$$\partial\Xi_0 := \{(2, 3), (2, 2), (2, 1), (3, 1), (4, 0)\}.$$

Moreover, $(\)_1(\)_2 b_\xi$ is in F iff $(\)_2(\)_1 b_\xi$ is. Hence, (5) also holds for the remaining $\xi \in \Xi$. This finishes the proof that, for this F and N , (b) of Theorem 1 holds.

It remains to show that, nevertheless, (a) of Theorem 1 does not hold. For this, observe that $(\)^{2,1}$ and $(\)^{4,0}$ are in $\ker N$, as is, e.g., $(\)_2 b_{1,6} = 7(\)^{1,7} + 3(\)^{3,3}$, hence $p = (\)^{1,7} + (\)^{3,3} + (\)^{5,0}$ is in the ideal generated by $\ker N$, making it impossible for N to be the restriction to $\Pi_1(F)$ of an ideal projector P with range F since this would place the nontrivial p in both $\ker P$ and $\text{ran } P$. ■

References

- [Bi] G. Birkhoff, *The algebra of multivariate interpolation*, in: Constructive approaches to mathematical models, C. V. Coffman and G. J. Fix (eds.), Academic Press, New York, 1979, 345–363.
- [Bo] C. de Boor, *Ideal interpolation*, in: Approximation Theory XI: Gatlinburg 2004, C. K. Chui, M. Neamtu, and L. L. Schumaker (eds.), Nashboro Press, Brentwood TN, 2005, 59–91.
- [CLO] David Cox, John Little, and Donal O’Shea, *Ideals, Varieties, and Algorithms*, Undergraduate Texts in Math., Springer-Verlag, New York, 1992.
- [Enc] *Encyclopädie der mathematischen Wissenschaften, 1. Band*, Teubner, Leipzig, 1898–1904.
- [E] L. Euler, *Calculus differentialis 1*, Ticini, Italy, 1787.
- [M] H. M. Möller, *Mehrdimensionale Hermite-Interpolation und numerische Integration*, Math. Z. 148 (1976) 107–118.
- [Mo] B. Mourrain, *A new criterion for normal form algorithms*, in: Applied Algebra, Algebraic Algorithms and Error-Correcting Codes (13th Intern. Symp., AAECC-13, Honolulu, Hawaii USA, Nov.’99, Proc.), Mark Fossorier, Hideki Imai, Shu Lin, Alan Pol (eds.), Springer Lecture Notes in Computer Science, 1719, Springer-Verlag, Heidelberg, 1999, 430–443.